

THE E-THEORETIC DESCENT FUNCTOR FOR GROUPOIDS

ALAN L. T. PATERSON

ABSTRACT. The paper establishes, for a wide class of locally compact groupoids Γ , the E-theoretic descent functor at the C^* -algebra level, in a way parallel to that established for locally compact groups by Guentner, Higson and Trout. The second section shows that Γ -actions on a $C_0(X)$ -algebra B , where X is the unit space of Γ , can be usefully formulated in terms of an action on the associated bundle B^\sharp . The third section shows that the functor $B \rightarrow C^*(\Gamma, B)$ is continuous and exact, and uses the disintegration theory of J. Renault. The last section establishes the existence of the descent functor under a very mild condition on Γ , the main technical difficulty involved being that of finding a Γ -algebra that plays the role of $C_b(T, B)^{cont}$ in the group case.

1. INTRODUCTION

In a number of situations, in particular for the assembly map, the Baum-Connes conjecture and index theory ([16, Theorem 3.4], [15, 5, 36, 37, 39] and many others) the descent homomorphism $j_G : KK_G(A, B) \rightarrow KK(C^*(G, A), C^*(G, B))$, where G is a locally compact group and A, B are G - C^* -algebras, is of great importance. (There is a corresponding result for the reduced crossed product algebras.) In non-commutative geometry, classical group symmetry does not suffice, and one requires smooth groupoids in place of Lie groups ([4, 5]), so that it is important to have available constructions, such as that which gives the descent homomorphism, for groupoid, rather than just for group, actions. To this end, the work of LeGall ([17, 7.2]) shows the existence of the KK_Γ -descent homomorphism for Γ -algebras, where Γ is a locally compact, σ -compact Hausdorff groupoid with left Haar system.

A similar issue arises when we consider E-theory rather than KK-theory. Non-equivariant E-theory is developed in [6, 5, 1]. Guentner, Higson and Trout gave a definitive account of group equivariant E-theory in their memoir [12]. In particular ([12, pp.47, 60ff.]), they established the group equivariant E-theoretic descent functor and used it in their definition of the E-theoretic assembly map. Another situation where the E-theoretic descent homomorphism is required is in the Bott periodicity theorem for infinite dimensional Euclidean space which was established by Higson, Kasparov and Trout ([14]), with its applications to the equivariant topological index and the Novikov higher signature conjecture. The descent homomorphism associates to an equivariant asymptotic morphism from A to B a canonical homomorphism from $K_G(A)$ to $K_G(B)$, and this is how it is used in [14]. The present paper studies the descent homomorphism in the much more general situation involving groupoids rather than groups. By modifying the method of [12],

Date: April, 2007.

1991 Mathematics Subject Classification. Primary: 19K35, 22A22, 46L80, 58B34.

Key words and phrases. groupoids, asymptotic morphism, descent functor.

we prove the existence of the groupoid descent homomorphism at the C^* -algebra level for a very wide class of groupoids.

We start by reformulating the concept of a Γ -action on a $C_0(X)$ -algebra. A Γ - C^* -algebra is defined in the literature ([17, 30]) as follows. Let X be the unit space of Γ and A be a $C_0(X)$ -algebra. Roughly, the latter means that A can be regarded as the $C_0(X)$ -algebra of continuous sections vanishing at infinity of a C^* -bundle A^\sharp of C^* -algebras A_x ($x \in X$). One pulls back A to Γ using the range and source maps r, s to obtain $C_0(\Gamma)$ -algebras r^*A, s^*A . An action of Γ on A is just a $C_0(\Gamma)$ -isomorphism $\alpha : s^*A \rightarrow r^*A$ for which the maps $\alpha_\gamma : A_{s(\gamma)} \rightarrow A_{r(\gamma)}$ compose in accordance with the rules for groupoid multiplication. It is desirable to have available an equivalent definition for a Γ -action along the lines of an action (in the usual sense) of a group on a C^* -algebra: in groupoid terms, this should involve a continuity condition for the map $\gamma \rightarrow \alpha_\gamma$ on the C^* -bundle A^\sharp . The specification of this continuity is very natural: we require that for each $a \in A$, the map $\gamma \rightarrow \alpha_\gamma(a_{s(\gamma)})$ be continuous. This definition is useful in a number of contexts, for example, in specifying the Γ -algebra of continuous elements in a C^* -algebra with an algebraic Γ -action, and in working with the covariant algebra $C^*(\Gamma, A)$.

The second section proves that the two definitions of Γ -action on A are equivalent. We survey the theory of C^* -bundles, in particular, the topologizing of A^\sharp . We require the well-known result, related to the Dauns-Hoffman theorem, that the “Gelfand transform” of A is an isomorphism onto $C_0(X, A^\sharp)$. A simple modification of a corresponding result by Dupré and Gillette ([9]) gives this result and we sketch it for completeness. (Another approach to this is given by Nilsen ([23]).)

Following the method of Guentner, Higson and Trout, we have to show that the functor $A \rightarrow C^*(\Gamma, A)$ is continuous and exact. This is proved in the third section. The continuity of this functor is proved in a way similar to that of the group case, while for exactness, we give a groupoid version of the corresponding theorem of N. C. Phillips ([28]) for locally compact groups. The proofs of these use the disintegration theorem of J. Renault.

The fourth section establishes our version of the descent homomorphism. The theory of groupoid equivariant asymptotic morphisms for Γ -algebras is developed using the work of R. Popescu ([30]), to whom I am grateful for helpful comments. In particular, in place of the non-equivariant $C_b(T, B)$, the $C_0(X)$ -algebra $C_b^X(T, B) = C_0(X)C_b(T, B)$ is used. A technical difficulty arises since (unlike the locally compact group case) there does not exist a natural algebraic Γ -action on $C_b^X(T, B)$. However, there is another natural bundle $C_b(T, B^\sharp) = \sqcup C_b(T, B_x)$ on which there is a simple algebraic Γ -action, derived from the given action on B , and a bundle map R from $C_b^X(T, B)^\sharp$ to $C_b(T, B^\sharp)$. We show that if Γ has local G -sets - a very mild, “transversal” condition satisfied by most groupoids that arise in practice - then we *can* find a canonical Γ -algebra $\mathfrak{B} \subset C_b^X(T, B)$ determined by the action of Γ on $C_b(T, B^\sharp)$. The map $B \rightarrow \mathfrak{B}$ is functorial and is the natural choice for defining the equivariant asymptotic algebra and the groupoid descent homomorphism.

2. GROUPOID C^* -ALGEBRAS

Let Γ be a locally compact, second countable, Hausdorff groupoid with a left Haar system λ . The unit space of Γ is denoted by X . The range and source maps $r, s : \Gamma \rightarrow X$ are given by: $r(\gamma) = \gamma\gamma^{-1}$, $s(\gamma) = \gamma^{-1}\gamma$. We now review Γ -spaces.

A Γ -space is a topological space M with an open, onto, projection map $p : M \rightarrow X$, $M_x = p^{-1}(x)$, and a continuous product map (a Γ -action) from $\Gamma \times_s M = \{(\gamma, m) : s(\gamma) = p(m)\} \rightarrow M$, so that the usual groupoid algebra holds: in particular, if $s(\gamma_2) = p(m)$, then $\gamma_1(\gamma_2 m) = (\gamma_1 \gamma_2)m$ whenever $s(\gamma_1) = r(\gamma_2)$, and $p(\gamma_2 m) = r(\gamma_2)$. In the case where each M_x is a C^* -algebra and each of the maps $z \rightarrow \gamma z$ is a $*$ -isomorphism from $M_{s(\gamma)}$ onto $M_{r(\gamma)}$, then we say that M is a Γ -space of C^* -algebras. We often write $\alpha_\gamma(z)$ in place of γz . For such an M , there is a natural groupoid $Iso(M)$ whose elements are the $*$ -isomorphisms t from some M_{x_1} onto M_{x_2} and with unit space X . Of course, $s(t) = x_1$ and $r(t) = x_2$ and the product is given by composition of maps. Then saying that $\gamma \rightarrow \alpha_\gamma$ is a Γ -action is equivalent to saying that the map is a groupoid homomorphism from Γ into $Iso(M)$. We call such an M a *continuous* Γ -space of C^* -algebras if the map $(\gamma, z) \rightarrow \alpha_\gamma(z)$ from $\Gamma \times_s M$ into M is continuous.

Let A be a separable C^* -algebra. We recall what it means for A to be $C_0(X)$ -algebra ([16, 2, 3, 9, 17, 23]). It means that there is a homomorphism θ from $C_0(X)$ into the center $ZM(A)$ of the multiplier algebra $M(A)$ of A such that $\theta(C_0(X))A = A$. A $C_0(X)$ -algebra A determines a family of C^* -algebras A_x ($x \in X$) where $A_x = A/(I_x A)$ with $I_x = \{f \in C_0(X) : f(x) = 0\}$. If J is a closed ideal of such an A , then the restriction map $f \rightarrow \theta(f)|_J$ makes J also into a $C_0(X)$ -algebra. Also, A/J is a $C_0(X)$ -algebra in the obvious way, and $(A/J)_x = A_x/J_x$.

A $C_0(X)$ -morphism from A to B , where A, B are $C_0(X)$ -algebras, is defined to be a $*$ -homomorphism $T : A \rightarrow B$ which is also a $C_0(X)$ -module map. In that case, T determines a $*$ -homomorphism $T_x : A_x \rightarrow B_x$ for each $x \in X$. The following discussion is very close to, but not quite contained, in the book on Banach bundles by Dupré and Gillette ([9]), and we will give a brief description of the modifications required. Some of the details will be needed later and do not seem to appear in the literature.

Let a_x be the image of $a \in A$ in A_x . Let $A^\# = \sqcup A_x$ and $p : A^\# \rightarrow X$ be the map: $p(a_x) = x$. Then (cf. [9, p.8]), $A^\#$ is a C^* -algebra family: the map $p : A^\# \rightarrow X$ is surjective, and each fiber $A_x = p^{-1}(x)$ is a C^* -algebra. Let \hat{a} be the section of $A^\#$ given by: $\hat{a}(x) = a_x$. By a C^* -family E being a C^* -bundle, we mean ([9, pp.6-9]) that E is a topological space with p open and continuous, that scalar multiplication, addition, multiplication and involution are continuous respectively from $\mathbb{C} \times E \rightarrow E$, from $E \times_X E \rightarrow E$, from $E \times_X E \rightarrow E$ and from $E \rightarrow E$, and the norm map $\|\cdot\| : E \rightarrow \mathbb{R}$ is upper semicontinuous and the following condition on the open sets for E holds: if W is open in E , $x \in X$ and the zero 0_x of E_x belongs to W , then there exists an $\epsilon > 0$ and an open neighborhood U of x such that

$$\{b \in p^{-1}(U) : \|b\| < \epsilon\} \subset W.$$

(Recall that a map $f : E \rightarrow \mathbb{R}$ is upper semicontinuous if, for each $e_0 \in E$ and each $\epsilon > 0$, there is an open neighborhood U of e_0 in E such that $f(e) < f(e_0) + \epsilon$ for all $e \in U$.)

We now discuss briefly how, in a natural way, $E = A^\#$ is a C^* -bundle, and the map $a \rightarrow \hat{a}$ is a $*$ -isomorphism from A onto $C_0(X, A^\#)$ (a ‘‘Gelfand’’ theorem) (cf. [17, 2.1.3], [23]). One proves first that each \hat{a} is upper semicontinuous. To this end, we modify the proof of the corresponding results ([9, Proposition 2.1, Corollary 2.2]) which are proved in [9] for the completely regular, rather than locally compact Hausdorff, case. The first of these results for our case can be stated as follows. For each $x \in X$, let \mathcal{N}_x be the family of relatively compact, open subsets V of X

containing x . For each $V \in \mathcal{N}_x$, let $f_V : X \rightarrow [0, 1]$ be continuous and such that it is 1 on a neighborhood of x in V and vanishes outside V . Then

$$(2.1) \quad \|\hat{a}(x)\| = \inf\{\|f_V a\| : V \in \mathcal{N}_x\}.$$

The upper semicontinuity of $\|\hat{a}\|$ follows since if $\epsilon > 0$ and V is chosen so that $\|\hat{a}(x)\| > \|f_V a\| - \epsilon$, then for y in a neighborhood of x , $\|\hat{a}(y)\| \leq \|f_V a\| < \|\hat{a}(x)\| + \epsilon$. (The equality (2.1) is due to J. Varela.)

The first part of the proof of (2.1) shows that $\|\hat{a}(x)\| \geq \inf\{\|f_V a\| : V \in \mathcal{N}_x\}$. This is the same as in the original Proposition 2.1. For the reverse inequality, let $\epsilon > 0$. Since A is a $C_0(X)$ -algebra, $a = fb$ for some $f \in C_0(X)$, $b \in A$, and using a bounded approximate identity in $C_0(X)$, there exists $F \in C_0(X)$ such that $0 \leq F \leq 1$, $F(x) = 1$ and $\|(1 - F)a\| < \epsilon$. As $(F - f_V)a \in I_x A$, and $a = f_V a + (1 - F)a + (F - f_V)a$,

$$\|\hat{a}(x)\| \leq \|f_V a + (1 - F)a\| \leq \|f_V a\| + \epsilon$$

and we obtain $\|\hat{a}(x)\| \leq \inf\{\|f_V a\| : V \in \mathcal{N}_x\}$.

Since we have (2.1), the conditions of ([9, Proposition 1.3]) (or of ([11, Proposition 1.6])) are satisfied, and A^\sharp is a C^* -bundle. A base for the topology on A^\sharp ([9, pp. 9-10, 16]) is given by sets of the form

$$(2.2) \quad U(a, \epsilon) = \{b_x \in A_x : x \in U, \|b_x - a_x\| < \epsilon\}$$

where $a \in A$, $\epsilon > 0$ and U is an open subset of X . Further, a local base at $z \in A_{x_0}$ is given by neighborhoods of the form $U(a, \epsilon)$ where a is any fixed element of A for which $a_{x_0} = z$, and \hat{A} is a closed $*$ -subalgebra of $C_0(X, A^\sharp)$. To see that $\hat{A} = C_0(X, A^\sharp)$, we just have to show (cf. ([9, Proposition 2.3])) that \hat{A} is dense in $C_0(X, A^\sharp)$. This follows by a simple partition of unity argument ([11, Proposition 1.7], [31, Lemma 5.3]). We then have the following theorem ([17, Théorème 2.1.1]). It is also proved by Nilsen ([23, Theorem 2.3]) who derives the Dauns-Hoffman theorem ([31, Theorem A.34]) from it.

Theorem 1. *With the above topology on A^\sharp , A^\sharp is a C^* -bundle over X . Further, the relative topology on each A_x is the norm topology. Last, the map $a \rightarrow \hat{a}$ is a $C_0(X)$ -isomorphism from A onto the $C_0(X)$ -algebra $C_0(X, A^\sharp)$ of continuous sections of A^\sharp that vanish at infinity.*

If A, B are $C_0(X)$ -algebras and $T : A \rightarrow B$ is a $C_0(X)$ -morphism, then $T^\sharp : A^\sharp \rightarrow B^\sharp$ is continuous, where $T^\sharp a_x = T_x(a_x) = (Ta)_x$. (In fact $(T^\sharp)^{-1}(U(Ta, \epsilon)) \supset U(a, \epsilon)$.)

Next, if \mathfrak{B} is a $C_0(X)$ -subalgebra of a $C_0(X)$ -algebra B then for any x , $I_x B \cap \mathfrak{B} = I_x \mathfrak{B}$ so that we can identify \mathfrak{B}^\sharp with a subbundle of B^\sharp , and the topology on \mathfrak{B}^\sharp is the relative topology.

We now recall how the (maximal) tensor product $A \otimes_{C_0(X)} B$ of two $C_0(X)$ -algebras is defined. For more information, see [2, 3, 17, 10]. One natural way to do this is to take $A \otimes_{C_0(X)} B$ to be the maximal $C_0(X)$ -balanced tensor product: so $A \otimes_{C_0(X)} B = (A \otimes_{\max} B)/I$, where I is the closed ideal generated by differences of the form $(af \otimes b - a \otimes fb)$ ($a \in A, b \in B, f \in C_0(X)$). The $C_0(X)$ -action on $A \otimes_{C_0(X)} B$ is determined by: $f(a \otimes b) = fa \otimes b = a \otimes fb$ for $f \in C_0(X)$. (Alternatively, one regards $A \otimes_{\max} B$ as a $C_0(X \times X)$ -algebra and “restricts to the diagonal”: $A \otimes_{C_0(X)} B = (A \otimes_{\max} B)/C_\Delta(A \otimes_{\max} B)$ where $C_\Delta = \{g \in C_0(X \times X) : g(x, x) = 0 \text{ for all } x \in X\}$.) Next, $(A \otimes_{C_0(X)} B)_x = A_x \otimes_{\max} B_x$.

If D is an ordinary C^* -algebra, then $D \otimes_{\max} B$ is a $C_0(X)$ -algebra in the natural way: $\theta(f)(d \otimes b) = d \otimes fb$. (Alternatively, one can identify the $C_0(X)$ -algebra $D \otimes_{\max} B$ with $(D \otimes_{\max} C_0(X)) \otimes_{C_0(X)} B$.) An important case of this is when $D = C_0(Z)$ (Z a locally compact Hausdorff space): then $C_0(Z, B) = C_0(Z) \otimes B$ is a $C_0(X)$ -algebra, and $C_0(Z, B)_x = C_0(Z) \otimes B_x = C_0(Z, B_x)$. It is easily checked that $(g \otimes b)_x = g \otimes b_x$, and it follows that for $F \in C_0(Z, B)$, $F_x(z) = F(z)_x \in B_x$.

Now let $B = C_0(Y)$ (Y a locally compact Hausdorff space) with the $C_0(X)$ -action on B given by a continuous map $q : Y \rightarrow X$: here $(fF)(y) = f(q(y))F(y)$ where $F \in C_0(Y)$, $f \in C_0(X)$. In this case, one writes $q^*A = A \otimes_{C_0(X)} C_0(Y)$. It is sometimes helpful to incorporate explicit mention of the map q in this tensor product by writing $A \otimes_{C_0(X), q} C_0(Y)$ in place of $A \otimes_{C_0(X)} C_0(Y)$. q^*A is actually also a $C_0(Y)$ -algebra in the obvious way: $(a \otimes F)F' = a \otimes FF'$ for $F, F' \in C_0(Y)$, and for each $y \in Y$, we have $(q^*A)_y = A_{q(y)}$. (The canonical map from $(q^*A)_y$ to $A_{q(y)}$ comes from sending $(a \otimes F)_y$ to $a_{q(y)}F(y)$.) Now let $Y \times_q A^\sharp = \{(y, a_{q(y)}) : y \in Y, a \in A\}$ with the relative topology inherited from $Y \times A^\sharp$. From the above, $Y \times_q A^\sharp$ is identified as a set with $(q^*A)^\sharp$. We now show that the spaces are homeomorphic when q is open.

Proposition 1. *If q is also open, then the identity map $i : Y \times_q A^\sharp \rightarrow (q^*A)^\sharp$ is a homeomorphism.*

Proof. A base for the topology of $(q^*A)^\sharp$ is given by sets of the form $W(a \otimes F, \epsilon)$ where W is a relatively compact open subset of Y , $F \in C_0(Y)$ is 1 on W and $a \in A$. Then $W(a \otimes F, \epsilon) = W \times_q q(W)(a, \epsilon)$ and the latter sets form a base for the topology of $Y \times_q A^\sharp$. \square

We now recall what is meant by a Γ -algebra A ([17, 30]). Form the balanced tensor products $s^*A = A \otimes_{C_0(X), s} C_0(\Gamma)$ and $r^*A = A \otimes_{C_0(X), r} C_0(\Gamma)$. From Theorem 1, $r^*A = C_0(\Gamma, (r^*A)^\sharp)$. Then A is called a Γ -algebra if there is given a $C_0(\Gamma)$ -isomorphism $\alpha : s^*A \rightarrow r^*A$ such that the induced isomorphisms $\alpha_\gamma : (s^*A)_\gamma \rightarrow (r^*A)_\gamma$ satisfy the groupoid multiplication properties: $\alpha_{\gamma\gamma'} = \alpha_\gamma \alpha_{\gamma'}$ whenever $r(\gamma') = s(\gamma)$ and $\alpha_{\gamma^{-1}} = (\alpha_\gamma)^{-1}$. Obviously, for each $x \in X$, α_x is the identity map on A_x .

As an example, suppose that Γ is a locally compact group G . Then $s^*A = r^*A = A \otimes C_0(G) = C_0(G, A)$. For $F \in C_0(G, A)$, we have $F_g = F(g)$ and by Proposition 1, $(C_0(G, A))^\sharp = G \times A$. If $\alpha : C_0(G, A) \rightarrow C_0(G, A)$ gives a G -action on $C_0(G, A)$, then since $(C_0(G, A))_g = A$, we get isomorphisms $\alpha_g : A \rightarrow A$. We are given that $\alpha_{gh} = \alpha_g \circ \alpha_h$ for all $g, h \in G$. Last since the map $g \rightarrow (\alpha(a \otimes k))_g = \alpha_g(a)k(g)$ belongs to $C_0(G, A)$, it follows that for each $a \in A$, the map $g \rightarrow \alpha_g(a)$ is norm continuous. So A is a G -algebra in the usual sense. (The converse is left to the reader.) We now show that the groupoid version of the preceding holds; a Γ - C^* -algebra A can then be viewed in terms of a Γ -action on A^\sharp . Note that the corollary characterizes Γ -action in terms exactly analogous to that of a group action.

Theorem 2. *A is a Γ -algebra if and only if A^\sharp is a continuous Γ -space of C^* -algebras A_x .*

Proof. Suppose that A is a Γ -algebra. So we are given a $C_0(\Gamma)$ -isomorphism $\alpha : s^*A \rightarrow r^*A$ with $\gamma \rightarrow \alpha_\gamma$ a homomorphism into $\text{Iso}(A^\sharp)$. Let $\beta = \alpha^\sharp$. Then the continuity of the map $(\gamma, z) \rightarrow \alpha_\gamma(z)$ follows, using Proposition 1, by composing

the following continuous maps:

$$\Gamma \times_s A^\# \xrightarrow{i} (s^*A)^\# \xrightarrow{\beta} (r^*A)^\# \xrightarrow{i^{-1}} \Gamma \times_r A^\# \xrightarrow{p_2} A^\#$$

where p_2 is the projection onto the second coordinate. So $A^\#$ is a continuous Γ -space of C^* -algebras.

Conversely, suppose that $A^\#$ is a continuous Γ -space of C^* -algebras. Define $\beta : \Gamma \times_s A^\# \rightarrow \Gamma \times_r A^\#$ by: $\beta(\gamma, z) = (\gamma, \alpha_\gamma(z))$. By assumption, β is continuous. The map β^{-1} is also continuous since it equals $(\text{inv} \otimes 1) \circ \beta \circ (\text{inv} \otimes 1)$, where $\text{inv}(\gamma) = \gamma^{-1}$. Then the map $F \rightarrow (i\beta i^{-1}) \circ F$ is a $C_0(\Gamma)$ -homomorphism from $C_0(\Gamma, (s^*A)^\#)$ into $C_0(\Gamma, (r^*A)^\#)$ which is an isomorphism since its inverse is the corresponding expression involving β^{-1} . By Theorem 1, this isomorphism determines a $C_0(\Gamma)$ -isomorphism $\alpha : s^*A \rightarrow r^*A$ and A is a Γ -algebra. \square

Corollary 1. *A is a Γ -algebra if and only if there is given a groupoid homomorphism $\gamma \rightarrow \alpha_\gamma$ from Γ into $\text{Iso}(A^\#)$ such that for each $a \in A$, the map $\gamma \rightarrow \alpha_\gamma(a_{s(\gamma)})$ is continuous.*

Proof. Suppose that we are given a groupoid homomorphism $\gamma \rightarrow \alpha_\gamma$ from Γ into $\text{Iso}(A^\#)$ such that for each $a \in A$, the map $\gamma \rightarrow \alpha_\gamma(a_{s(\gamma)})$ is continuous. Let $\{(\gamma_\delta, z_\delta)\}$ be a net in $\Gamma \times_s A^\#$ converging to some (γ_0, z_0) . We show that $\alpha_{\gamma_\delta}(z_\delta) \rightarrow \alpha_{\gamma_0}(z_0)$. Let $a, c \in A$ be such that $z_0 = a_{s(\gamma_0)}$ and $\alpha_{\gamma_0}(z_0) = c_{r(\gamma_0)}$. Let $V(c, \epsilon)$ be a neighborhood of $c_{r(\gamma_0)}$ in $A^\#$. By continuity of the map $\gamma \rightarrow \alpha_\gamma(a_{s(\gamma)})$, there exists an open neighborhood Z of γ_0 in Γ and a δ_1 such that for all $\delta \geq \delta_1$, $\alpha_\gamma(a_{s(\gamma)}) \in V(c, \epsilon/2)$ for all $\gamma \in Z$. Since $z_\delta \rightarrow z_0$, we can also arrange that $z_\delta \in s(Z)(a, \epsilon/2)$ for all $\delta \geq \delta_1$. So for all $\delta \geq \delta_1$, $\|z_\delta - a_{s(\gamma_\delta)}\| < \epsilon/2$, giving $\|\alpha_{\gamma_\delta}(z_\delta) - \alpha_{\gamma_\delta}(a_{s(\gamma_\delta)})\| < \epsilon/2$. Since $\alpha_{\gamma_\delta}(a_{s(\gamma_\delta)}) \rightarrow c_{r(\gamma_0)}$, we can also suppose that for all $\delta \geq \delta_1$, $\alpha_{\gamma_\delta}(a_{s(\gamma_\delta)}) \in V(c, \epsilon/2)$, i.e. $\|\alpha_{\gamma_\delta}(a_{s(\gamma_\delta)}) - c_{r(\gamma_\delta)}\| < \epsilon/2$. By the triangular inequality, $\alpha_{\gamma_\delta}(z_\delta) \in V(c, \epsilon)$ ($\delta \geq \delta_1$), and $\alpha_{\gamma_\delta}(z_\delta) \rightarrow \alpha_{\gamma_0}(z_0)$. By Theorem 2, A is a Γ -algebra. The converse also follows from Theorem 2. \square

Now suppose that A is a Γ -algebra and J is a closed ideal of A that is a Γ -subalgebra of A in the natural way, i.e. for each $\gamma \in \Gamma$, $j \in J$, we have $\alpha_\gamma(j_{s(\gamma)}) \in J_{r(\gamma)}$. Using the continuity of the canonical map from $A^\#$ to $(A/J)^\#$ and Corollary 1, it is easy to prove that the $C_0(X)$ -algebra A/J is also a Γ -algebra in the natural way, and we have a short exact sequence of Γ -algebras:

$$(2.3) \quad 0 \rightarrow J \rightarrow A \rightarrow A/J \rightarrow 0.$$

Next suppose that A is a $C_0(X)$ -algebra, and that the A_x 's form a Γ -space of C^* -algebras. So we can say that A has an *algebraic* Γ -action (with no continuity condition on the maps $\gamma \rightarrow \alpha_\gamma(a_{s(\gamma)})$). We wish to define a C^* -subalgebra A^{cont} of A on which the Γ -action is continuous. For this result, we require that Γ have *local $r - G$ -sets* (cf. [32, p.10], [25, p.44]). This means that for each $\gamma_0 \in \Gamma$, there exists an open neighborhood U of $r(\gamma_0)$ in X and a subset W of Γ containing γ_0 such that $r_W = r|_W$ is a homeomorphism from W onto U . Most locally compact groupoids that arise in practice have local $r - G$ -sets (e.g. Lie groupoids, r -discrete groupoids and transformation group groupoids).

Proposition 2. *Let Γ have local $r - G$ -sets, A have an algebraic Γ -action and define*

$$A^{cont} = \{a \in A : \text{the map } \gamma \rightarrow \alpha_\gamma(a_{s(\gamma)}) \text{ is continuous}\}.$$

Then A^{cont} is a Γ -subalgebra of A .

Proof. It is obvious from the definition of a C^* -bundle that A^{cont} is a $*$ -subalgebra of A . Let $a_n \rightarrow a$ in A with $a_n \in A^{cont}$ for all n . Then $\alpha_\gamma((a_n)_{s(\gamma)}) \rightarrow \alpha_\gamma(a_{s(\gamma)})$ uniformly in γ . Adapting the proof of the elementary result that a uniform limit of continuous functions is continuous - one uses also the upper semicontinuity of the norm on $A^\#$ - it follows that the map $\gamma \rightarrow \alpha_\gamma(a_{s(\gamma)})$ is continuous, i.e. $a \in A^{cont}$. So A^{cont} is a C^* -subalgebra of A . Next, if $f \in C_0(X)$, then $\alpha_\gamma((fa)_{s(\gamma)}) = f(s(\gamma))\alpha_\gamma(a_{s(\gamma)})$, and the map $\gamma \rightarrow \alpha_\gamma((fa)_{s(\gamma)})$ is continuous. So $C_0(X)A^{cont} \subset A^{cont}$. Also, if $a \in A^{cont}$, then $a = f'a'$ for some $f' \in C_0(X)$, $a' \in A$, and so $a = \lim e_n(f'a')$ where $\{e_n\}$ is a bounded approximate identity for $C_0(X)$. So $C_0(X)A^{cont} = A^{cont}$, and A^{cont} is a $C_0(X)$ -algebra. Last, we have to show that if $a \in A^{cont}$ and $\gamma_0 \in \Gamma$, then $\alpha_{\gamma_0}(a_{s(\gamma_0)}) \in (A^{cont})_{r(\gamma_0)}$. Let W, U be as above so that $r_W : W \rightarrow U$ is a homeomorphism. Let $f \in C_c(U)$ be such that $f(r(\gamma_0)) = 1$. Then the section g of $A^\#$ given by: $g(u) = f(u)\alpha_{r_W^{-1}(u)}(a_{s(r_W^{-1}(u))})$ belongs to $C_c(X, A^\#)$. By Theorem 1, there exists $b \in A$ such that $b_u = g(u)$ for all $u \in X$. Since $\alpha_\gamma(b_{s(\gamma)}) = f(s(\gamma))\alpha_{\gamma \cdot r_W^{-1}(s(\gamma))}(a_{s(\gamma \cdot r_W^{-1}(s(\gamma)))})$ and $a \in A^{cont}$, we see that $b \in A^{cont}$. \square

Now let A, B be Γ -algebras. The tensor product $A \otimes_{C_0(X)} B$ is a Γ -algebra ([17, 3.1.2]) in a natural way. Indeed, using the associativity of the balanced tensor product ([2, p.90]) and the equality $C_0(\Gamma) \otimes_{C_0(\Gamma)} C_0(\Gamma) = C_0(\Gamma)$, we obtain $q^*(A \otimes_{C_0(X)} B) = q^*A \otimes_{C_0(\Gamma)} q^*B$ ($q = s, r$). The Γ -action on $A \otimes_{C_0(X)} B$ is then given by $\alpha \otimes \beta$, where α, β are the Γ -actions on A, B . Further $(\alpha \otimes \beta)\gamma = \alpha_\gamma \otimes \beta_\gamma$ (recalling that $(A \otimes_{C_0(X)} B)_x = A_x \otimes_{max} B_x$). Also, if A is just a C^* -algebra and B is a Γ -algebra, then the $C_0(X)$ -algebra $A \otimes_{max} B$ is a Γ -algebra: we identify $q^*(A \otimes_{max} B)$ with $A \otimes_{max} q^*B$ with $q = s, r$, and the Γ -action is given by $I \otimes \beta$. (Alternatively, one can reduce this to the earlier case by using $q^*((A \otimes C_0(X)) \otimes_{C_0(X)} B)$.) In particular, if B is a Γ -algebra, then the $C_0(X)$ -algebra $C_0(T, B)$ is also a Γ -algebra, and the action is given by:

$$(2.4) \quad \alpha_\gamma(F_{s(\gamma)})(t) = \alpha_\gamma(F(t)_{s(\gamma)}).$$

A Γ -homomorphism ([17, Definition 3.1.2]) from A to B is a $C_0(X)$ -homomorphism $\phi : A \rightarrow B$ such that for all $\gamma \in \Gamma$,

$$(2.5) \quad \phi_{r(\gamma)}\alpha_\gamma = \beta_\gamma\phi_{s(\gamma)}$$

where α, β denote respectively the actions of Γ on A and B . It is simple to check that with Γ -homomorphisms as morphisms, the class of Γ -algebras forms a category.

3. CONTINUITY AND EXACTNESS

Next, we need the notion of a *crossed product* of Γ by a Γ -algebra A ([33], [17]). We will need to use the profound disintegration theorem of J. Renault of [33]. Renault develops a groupoid version of the theory of twisted covariance algebras for locally compact groups, and working in a very general context, constructs a C^* -algebra $C^*(\Gamma, \Sigma, A, \lambda)$ where A is a Γ -algebra and S is a bundle of abelian

groups over X with Γ acting on the fibers and Σ is a groupoid given by an exact sequence of groupoids. (Also used in the construction is a homomorphism χ on S that we don't need to go into for our present purposes.) For the special case of the groupoid crossed product, we take $S = X = \Gamma^0$ and $\Sigma = \Gamma$. In that context, we put on the algebra $r_c^*A = C_c(\Gamma, (r^*A)^\#) \cong C_c(\Gamma)r^*A$, a product and involution given by:

(3.1)

$$F_1 * F_2(\gamma) = \int_{\Gamma^{r(\gamma)}} F_1(\gamma') \alpha_{\gamma'}(F_2(\gamma'^{-1}\gamma)) d\lambda^{r(\gamma)}(\gamma') \quad (F_1)^*(\gamma) = \alpha_\gamma(F_1(\gamma^{-1})^*).$$

(The proof that $F_1 * F_2 \in r_c^*A$ is given by P.-Y. Le Gall in [17, Proposition 7.1.1].) Next, r_c^*A is a normed $*$ -algebra with isometric involution under the I -norm $\|\cdot\|_I$, where

$$\|F\|_I = \max\{\|F\|_r, \|F^*\|_r\}$$

and

$$\|F\|_r = \sup_{x \in X} \int_{\Gamma^x} \|F(\gamma)\| d\lambda^x(\gamma).$$

The enveloping C^* -algebra of $(r_c^*A, \|\cdot\|_I)$ is then defined to be the crossed product $C^*(\Gamma, A)$.

A very simple example of this is provided by the case where $A = C_0(X)$ with the usual action of Γ on X : $\alpha_\gamma(s(\gamma)) = r(\gamma)$. In that case, as is easily checked, $A^\# = X \times \mathbb{C}$, $\alpha_\gamma : \mathbb{C}_{s(\gamma)} \rightarrow \mathbb{C}_{r(\gamma)}$ is the identity map, $s^*A = r^*A = C_0(\Gamma)$, and $\alpha : C_0(\Gamma) \rightarrow C_0(\Gamma)$ the identity map. Of course, $(r^*A)^\#$ is just $\Gamma \times \mathbb{C}$, and $r_c^*A = C_c(\Gamma)$. Then $C^*(\Gamma, A)$ is just the C^* -algebra $C^*(\Gamma)$ of the groupoid ([32]).

We now turn to Renault's disintegration theorem for representations of $C^*(\Gamma, A)$ - for a detailed exposition for the case $C^*(\Gamma)$ see [20]. The theory uses the fundamental papers of Ramsay ([34, 35]). We first formulate [33, Lemme 4.5] in $C_0(X)$ -algebra terms. Let A be a $C_0(X)$ -algebra, $\mathfrak{H} = \{H_x\}_{x \in X}$ a Hilbert bundle and μ a probability measure on X . Let $H = L^2(X, \mu, \mathfrak{H})$. We will say that a non-degenerate representation $\pi : A \rightarrow B(H)$ is a $C_0(X)$ -representation (for (X, μ, \mathfrak{H})) if π commutes with the $C_0(X)$ -actions on A and $B(H)$, i.e. for all $f \in C_0(X)$ and all $a \in A$, $T_f \pi(a) = \pi(a) T_f = \pi(fa)$, where T_f is the multiplication operator on $L^2(H)$ associated with f . By taking strong operator limits, we get that every a commutes with every T_f for $f \in L^\infty(X, \mu)$, i.e. with every diagonalizable operator. So ([7, II, 2, 5, Corollary]) every $\pi(a)$ is decomposable, and from [8, Lemma 8.3.1], π is a direct integral of representations π_x of A . Further, for each $f \in C_0(X)$, $\pi_x(fa) = f(x)\pi_x(a)$ so that π_x is a representation of A_x on H_x . The π_x 's are non-degenerate a. e. by [8, 8.1.5]. We now discuss what is meant by a covariant representation of (Γ, A) .

Let μ be quasi-invariant on X , $\nu = \int_X \lambda^x d\mu(x)$ ([32, pp.22-23], [25, Ch. 3]): quasi-invariance means that $\nu \sim \nu^{-1}$. Let U be a Borel subset of X which is μ -conull. Then $\Gamma|_U = r^{-1}(U) \cap s^{-1}(U)$ is a Borel groupoid which is ν -conull in Γ . Then $\Gamma|_U$ equipped with the restrictions of μ, ν to $U, \Gamma|_U$ is a measured groupoid, called the *inessential contraction* of Γ to U . Next we are given a Hilbert bundle $\mathfrak{H} = \{H_x\}_{x \in X}$; $Iso(X * \mathfrak{H})$ is the Borel groupoid of unitaries $U_{y,x} : H_x \rightarrow H_y$ as x, y range over X . (See [20, Chapter 3].) A *covariant representation* (or a *representation of the dynamical system* (Γ, Γ, A) in the terminology of ([33, p.79])) (L, π) of the pair (Γ, A) consists of:

- (i) a Borel homomorphism $L : \Gamma|_U \rightarrow Iso(X * \mathfrak{H})|_U$,

- (ii) a (non-degenerate) $C_0(X)$ -representation $\pi = \int^\oplus \pi_x d\mu(x)$ of A on $H = L^2(X, \mu, \mathfrak{H})$: for each $a \in A$,

$$\pi(a) = \int^\oplus \pi_x(a_x) d\mu(x),$$

- (iii) for all $\gamma \in \Gamma|_U$ and $a \in A$, we have

$$(3.2) \quad L\gamma\pi_s(\gamma)(a_s(\gamma))L\gamma^{-1} = \pi_r(\gamma)(\alpha_\gamma(a_s(\gamma))).$$

A. Ramsay ([34, 35, 20]) showed, at least in the case $A = C_0(X)$ above we can actually take $U = X$. However, because of the conullity of U , we can effectively regard the pair (L, π) as defined on Γ rather than $\Gamma|_U$ and will usually leave the U implicit.

Every covariant representation (L, π) of Γ integrates up to give a representation Φ of $C^*(\Gamma, A)$. Indeed, from [33, p.80], for $F \in r_c^*A$ and $\xi, \eta \in \mathfrak{H}$,

$$(3.3) \quad \langle \Phi(F)\xi, \eta \rangle = \int \langle \pi_r(\gamma)(F\gamma)L\gamma\xi_s(\gamma), \eta_r(\gamma) \rangle d\nu_0(\gamma)$$

where, as usual ([32, p.52], [25, 3.1]) $d\nu_0 = D^{-1/2} d\nu$ with $D = d\nu/d\nu^{-1}$.

Conversely, every representation Ψ of $C^*(\Gamma, A)$ on a Hilbert space K is equivalent to such a Φ . Indeed, from [33, p.88], elements ϕ, h of the algebras $C_c(\Gamma), C_c(X, A^\#)$ act as left multipliers on r_c^*A where:

$$(3.4) \quad \phi * F(\gamma) = \int_{\Gamma^{r(\gamma)}} \phi(\gamma')\alpha_{\gamma'}(F(\gamma'^{-1}\gamma))\lambda^{r(\gamma)}(\gamma'), \quad (hF)(\gamma) = (h \circ r)(\gamma)F(\gamma).$$

Renault shows that there are representations L', π' of $C_c(\Gamma), C_c(X, A^\#) = C_c(X)A$ on K and determined by:

$$(3.5) \quad \Psi(\phi * F) = L'(\phi)\Psi(F), \quad \Psi(hF) = \pi'(h)\Psi(F).$$

Renault first studies the representation L' of $C_c(\Gamma)$ and obtains a quasi-invariant measure μ on X and a measurable Hilbert bundle $\mathfrak{H} = \{H_x\}$ over X such that K can be identified with $L^2(X, \mu; \mathfrak{H})$. Then L' is disintegrated into a representation L of the groupoid Γ , and the $\pi'(a)$'s are decomposable on \mathfrak{H} : $\pi'(a) = \{\pi_x(a_x)\}$ (with π in place of π'). He then shows that L, π can be taken to be such that the pair (L, π) is a covariant pair whose integrated form is equivalent to Ψ .

We now discuss exactness and continuity for functors. So let \mathbf{F} be a functor from the category of Γ -algebras with Γ -homomorphisms as morphisms into the category of ordinary C^* -algebras with $*$ -homomorphisms as morphisms. Following [12, p.19, ff.], we say that \mathbf{F} is *exact* if for every short exact sequence of Γ -algebras

$$0 \rightarrow J \rightarrow A \rightarrow A/J \rightarrow 0$$

the induced sequence of ordinary C^* -algebras

$$0 \rightarrow \mathbf{F}(J) \rightarrow \mathbf{F}(A) \rightarrow \mathbf{F}(A/J) \rightarrow 0$$

is exact. Now let I be a closed interval $[a, b]$, B be a Γ -algebra and IB be the Γ -algebra $C(I) \otimes B = C(I, B)$. For each $k \in \mathbf{F}(IB)$, we can associate a function $\hat{k} : I \rightarrow \mathbf{F}(B)$ by setting $\hat{k}(t_0) = \mathbf{F}(ev_{t_0})(k)$ where $ev_{t_0} : IB \rightarrow B$ is evaluation at t_0 : $ev_{t_0}(g) = g(t_0)$. (Note that ev_{t_0} is a Γ -homomorphism.) The functor \mathbf{F} is said to be *continuous* if every \hat{k} is continuous. The map $k \rightarrow \hat{k}$ then gives a homomorphism from $\mathbf{F}(IB)$ into $I\mathbf{F}(B)$. Later, we will need to replace the finite interval I in $C(I, B)$ by the infinite interval T . We cannot replace $C(I, B)$ by $C_b(T, B)$ since

the latter does not admit a Γ -action in any natural way. However, the theory can be made to work, as we will see later (Proposition 5) by replacing $C_b(T, B)$ by a $C_0(X)$ -subalgebra \mathfrak{B} with a covering Γ -action.

An exact, continuous functor \mathbf{F} will now be constructed from the category of Γ -algebras with Γ -homomorphisms as morphisms into the category of ordinary C^* -algebras with $*$ -homomorphisms as morphisms. For a Γ -algebra A , define $\mathbf{F}(A) = C^*(\Gamma, A)$. We need to specify what \mathbf{F} does to morphisms. Let B also be a Γ -algebra and $\phi : A \rightarrow B$ be a Γ -homomorphism. Define, for each $F \in r_c^* A$, a section $\tilde{\phi}(F) : \Gamma \rightarrow (r^* B)^\#$ by:

$$(3.6) \quad \tilde{\phi}(F)(\gamma) = \phi_{r(\gamma)}(F_\gamma).$$

We note that $\tilde{\phi}(F)$ is just the same as $(r^* \phi)(F) \in r_c^* B = C_c(\Gamma, (r^* B)^\#)$. Using (3.1) and (2.5), we obtain that for $F_1, F_2 \in r_c^* A$ and each $\gamma \in \Gamma$,

$$(3.7) \quad \tilde{\phi}(F_1 * F_2)(\gamma) = \phi_{r(\gamma)}((F_1 * F_2)(\gamma)) = (\tilde{\phi}(F_1) * \tilde{\phi}(F_2))(\gamma)$$

and $\tilde{\phi}((F_1)^*) = (\tilde{\phi}(F_1))^*$. So $\tilde{\phi}$ is a $*$ -homomorphism from $r_c^* A$ to $r_c^* B$. It is continuous for the respective C^* -norms since $\|\tilde{\phi}(F)\|_I \leq \|F\|_I$ so that $\pi \circ \tilde{\phi}$ is a representation of $C^*(\Gamma, A)$ whenever π is a representation of $C^*(\Gamma, B)$. We set $\mathbf{F}(\phi) = \tilde{\phi}$. It is easy to check that \mathbf{F} is a functor. In the following, $A \otimes B$ means $A \otimes_{\max} B$.

Theorem 3. *The functor \mathbf{F} is continuous and exact.*

Proof. (a) We first show the continuity of F (cf. [12, Lemma 4.11] where the locally compact group case is sketched). Let A be an ordinary C^* -algebra and B be a Γ - C^* -algebra and recall that $A \otimes B$ is a Γ -algebra. We show that

$$(3.8) \quad A \otimes C^*(\Gamma, B) \cong C^*(\Gamma, A \otimes B).$$

There is a natural $C_0(\Gamma)$ -isomorphism Φ from $A \otimes r^* B$ onto $r^*(A \otimes B)$ determined by: $\Phi(a \otimes (b \otimes_{C_0(X)} F)) = (a \otimes b) \otimes_{C_0(X)} F$ (e.g. [30, Corollary 1.3]). The map Φ restricts to an isomorphism, also denoted Φ , from $A \otimes_{\text{alg}} r_c^* B$ onto a subalgebra of $r_c^*(A \otimes B)$. Φ is also an isomorphism when $A \otimes_{\text{alg}} r_c^* B, r_c^*(A \otimes B)$ are given the convolution product and involution of (3.1). Give $A \otimes_{\text{alg}} r_c^* B, r_c^*(A \otimes B)$ the norms that they inherit as (dense) subalgebras of $A \otimes C^*(\Gamma, B), C^*(\Gamma, A \otimes B)$. We note that $\Phi(A \otimes r_c^* B)$ is $\|\cdot\|_I$ -dense in $r_c^*(A \otimes B)$ and so also dense in $C^*(\Gamma, A \otimes B)$. We show that Φ is isometric.

A representation π' of $r_c^*(A \otimes B)$ is determined by a covariant pair (L, π) where π is a representation of $A \otimes B$ on some $H = L^2(X, \mu, \mathfrak{H})$. Then ([22, Theorem 6.3.5]) there exist non-degenerate, commuting representations π_1, π_2 of A, B on H such that $\pi(a \otimes b) = \pi_1(a)\pi_2(b)$. Further, using bounded approximate identities in A, B , π_2 is a $C_0(X)$ -representation and π_1 commutes with the $C_0(X)$ -multiplication operators T_f on H . Disintegrating, we get

$$\pi_1(a)_x(\pi_2)_x(b_x) = \pi_x(a \otimes b_x) = (\pi_2)_x(b_x)\pi_1(a)_x$$

almost everywhere, and $L_\gamma \pi_1(a)_{s(\gamma)}(\pi_2)_{s(\gamma)}(b_{s(\gamma)})L_{\gamma^{-1}} = \pi_1(a)_{r(\gamma)}(\pi_2)_{r(\gamma)}(b_{r(\gamma)})$. It follows that (L, π_2) is covariant for B and $L_\gamma \pi_1(a)_{s(\gamma)}L_{\gamma^{-1}} = \pi_1(a)_{r(\gamma)}$ a.e.. Let Φ_2 be the integrated form of (L, π_2) . Then from (3.3) and the above, the representations π_1, Φ_2 commute, and so the C^* -semi-norm that they induce on $A \otimes_{\text{alg}} C^*(\Gamma, B)$ is \leq the maximum tensor product norm. Since $\pi'(\Phi(w)) = (\pi_1 \otimes \Phi_2)(w)$ ($w \in A \otimes_{\text{alg}} r_c^* B$), it follows that $\|\Phi(w)\| \leq \|w\|$.

On the other hand, each representation π of $A \otimes C^*(\Gamma, B)$ is determined by a pair of commuting representations π_1, π_2 of $A, C^*(\Gamma, B)$ on some H . Then π_2 disintegrates into a covariant pair (π'_2, L) and we can identify $H = L^2(X, \mu, \mathfrak{H})$. Using (3.5), π_1 and π'_2 commute and L' and π_1 commute. Also, π_1 commutes with the T_f 's ($f \in C_0(X)$). From the proof of the disintegration theorem, $L\gamma\pi_1(a)_{s(\gamma)}L\gamma^{-1} = \pi_1(a)_{r(\gamma)}$ almost everywhere. Then $(\pi_1 \otimes \pi'_2, L)$ is a covariant representation for $A \otimes B$ and so determines a representation ϕ of $C^*(\Gamma, A \otimes B)$. Then on the range P of Φ , $\pi \circ \Phi^{-1} = \phi$, and it follows that $\|\Phi^{-1}(z)\| \leq \|z\|$ for all $z \in P$. So Φ is isometric, and so extends to an isomorphism from $A \otimes C^*(\Gamma, B)$ onto $C^*(\Gamma, A \otimes B)$, giving (3.8). For the continuity of \mathbf{F} , we take $A = C(I)$ where I is some $[a, b]$. Then using (3.8), let $k \in \mathbf{F}(IB) = C^*(\Gamma, IB) \cong C(I, C^*(\Gamma, B))$. When k belongs (under the isomorphism Φ) to the dense subalgebra $C(I) \otimes_{alg} r_c^* B$ of $C^*(\Gamma, IB)$, we use (3.6) to show that $\hat{k}(t_0) = \widehat{ev_{t_0}}(k) = k(t_0)$. By uniform convergence in $C(I, C^*(\Gamma, B))$, the same is true for $k \in C^*(\Gamma, IB)$, and \mathbf{F} is continuous.

(b) For exactness, one modifies the proof by N. C. Phillips of the corresponding result for the group case ([28, Lemma 2.8.2]). Let

$$0 \rightarrow J \xrightarrow{\chi} A \xrightarrow{\phi} B \rightarrow 0$$

be a Γ -equivariant short exact sequence of Γ -algebras. With $j = \mathbf{F}(\chi), \psi = \mathbf{F}(\phi)$, we have to show that

$$0 \rightarrow C^*(\Gamma, J) \xrightarrow{j} C^*(\Gamma, A) \xrightarrow{\psi} C^*(\Gamma, B) \rightarrow 0$$

is a short exact sequence of C^* -algebras. So we have to show that (1) j is injective, (2) $\psi \circ j = 0$, (3) $\ker \psi \subset j(C^*(\Gamma, J))$, and (4) ψ is surjective.

(1) Let Φ be a representation of $C^*(\Gamma, J)$. Let (L, π) be a disintegration of Φ on $H = L^2(X, \mu; \mathfrak{H})$ as earlier. Regarding the elements of A as multipliers on J in the obvious way, π extends to homomorphism π' of A . Further,

$$(3.9) \quad \pi'(a)\xi = \lim \pi(ae_n)\xi$$

where $\{e_n\}$ is sequence that is a bounded approximate identity for J . It follows that every $\pi'(a)$ is decomposable, and there is, for each $x \in X$, a representation π'_x of A_x on H_x such that for each $a \in A$,

$$\pi'(a) = \int^\oplus \pi'_x(a_x) d\mu(x).$$

Further, for a. e. x , π'_x is non-degenerate. Next, the restriction of π' to J is just π and so by the uniqueness a. e. of the decomposition of a decomposable operator ([7, II, 2, 3, Corollary]) and after removing a null set from U , we can suppose that π'_u restricts to π_u for all $u \in U$. Then $\pi'_u(a_u) = \lim \pi_u(a_u(e_n)_u)$ in the strong operator topology. We claim that the $\{\pi'_u\}$ are covariant for the $L\gamma$'s. Indeed, let $a \in A$. Then with convergence in the strong operator topology,

$L\gamma\pi'(a_{s(\gamma)})L\gamma^{-1} = \lim L\gamma\pi_{s(\gamma)}(a_{s(\gamma)}(e_n)_{s(\gamma)})L\gamma^{-1} = \lim \pi_{r(\gamma)}(\alpha\gamma(a_{s(\gamma)}(e_n)_{s(\gamma)}))$
 $= \lim \pi_{r(\gamma)}(\alpha\gamma(a_{s(\gamma)}))(\alpha\gamma((e_n)_{s(\gamma)})) = \pi'_{r(\gamma)}(\alpha\gamma(a_{s(\gamma)}))$. (Here we use the fact that $\{\alpha\gamma((e_n)_{s(\gamma)})\}$ is a bounded approximate identity for $J_{r(\gamma)}$.) So the pair (L, π') is a covariant representation of (Γ, A) and its integrated form Φ' is a representation of $C^*(\Gamma, A)$. Further, since χ is the identity map, $\Phi(g) = \Phi'(j(g))$ for all $g \in r_c^* J \subset r_c^* A$. It follows that $\|g\| \leq \|j(g)\|$ for all $g \in r_c^* J$, and by the continuity of j , this inequality extends to $C^*(\Gamma, J)$, and j is injective.

(2) $\phi \circ \chi = 0$, $\mathbf{F}(0) = 0$ and \mathbf{F} is a functor.

(3) From (1), j identifies $C^*(\Gamma, J)$ with a closed ideal of $C^*(\Gamma, A)$. Let $g_0 \in C^*(\Gamma, A)$ and suppose that $g_0 \notin C^*(\Gamma, J)$. Then there exists a representation Φ of $C^*(\Gamma, A)$ such that Φ annihilates $C^*(\Gamma, J)$ while $\Phi(g_0) \neq 0$. Let (L, π) be the covariant representation of (Γ, A) associated with Φ . If $h \in C_c(X, J^\sharp)$, $F \in r_c^*A$, then $hF \in r_c^*J$, and so by (3.5), $\pi(h) = 0$. So $\pi|_J = 0$ and π determines a $C_0(X)$ -representation π_1 of $A/J = B$. Also for $a \in A$, since $\pi_1 \circ \phi = \pi$,

$$(3.10) \quad (\pi_1)_x(\phi_x(a_x)) = \pi_x(a_x),$$

It is easy to check that the pair (L, π_1) is covariant for (Γ, B) . Let Φ_1 be the representation of $C^*(\Gamma, B)$ that is the integrated form of (L, π_1) . A simple calculation using (3.3) and (3.10) gives $\Phi = \Phi_1 \circ \psi$. Since $\Phi(g_0) \neq 0$, we must have $\psi(g_0) \neq 0$. So $\ker \psi \subset C^*(\Gamma, J)$.

(4) Let $F \in r_c^*B$, $\epsilon > 0$. Then there exist $F_i \in C_c(\Gamma)$, $b_i \in B$ such that $\|\sum_i b_i \otimes_{C_0(X)} F_i - F\| < \epsilon$ in r_c^*B . By multiplying by a fixed function $g \in C_c(\Gamma)$ with $g = 1$ on the support C of F , we can suppose that there is a fixed compact set K , independent of ϵ , containing the supports of F and the F_i 's. Then $\|\sum_i b_i \otimes_{C_0(X)} F_i - F\|_I \leq \epsilon \sup_x \lambda^x(K \cup K^{-1})$. Since $\sum_i b_i \otimes_{C_0(X)} F_i \in \psi(r_c^*A)$, it follows that ψ is surjective. \square

For later use, in the argument of (a) above, we can take in (3.8) $A = C_0(T)$ to obtain that for $k \in C_0(T, B)$, the function

$$(3.11) \quad \hat{k} \in C_0(T, \mathbf{F}(B)).$$

4. THE DESCENT HOMOMORPHISM

The theory of Γ -equivariant asymptotic homomorphisms was developed by R. Popescu ([30]). (The case where Γ is a locally compact group was treated in [12].) Recall first that in the non-equivariant case, one is given two C^* -algebras A, B . Let $T = [1, \infty)$. One defines $\mathfrak{A}B = C_b(T, B)/C_0(T, B)$. (The algebras $\mathfrak{A}^n B$ ($n \geq 2$) are defined inductively: $\mathfrak{A}^n B = \mathfrak{A}(\mathfrak{A}^{(n-1)}B)$, but for convenience, we restrict our discussion to the case $n = 1$.) An asymptotic morphism is a $*$ -homomorphism ϕ from A into $\mathfrak{A}B$.

The theory of asymptotic morphisms in the $C_0(X)$ -category requires natural and simple modifications ([24, 30]). The algebras A, B are, of course, taken to be $C_0(X)$ -algebras. However $C_b(T, B)$ is not a $C_0(X)$ -algebra under the natural homomorphism $\theta : C_0(X) \rightarrow ZM(C_b(T, B))$, where $(\theta(f)F)(t) = fF(t)$ ($f \in C_0(X)$, $F \in C_b(T, B)$). The reason is that $C_0(X)C_b(T, B) \neq C_b(T, B)$. Instead, one replaces $C_b(T, B)$ by its submodule $C_b^X(T, B) = C_0(X)C_b(T, B)$ which is a $C_0(X)$ -algebra. To ease the notation, we write $C_b(T, B)$ instead of $C_0(X)C_b(T, B)$ when no misunderstanding can arise. Recall (earlier) that $C_0(T, B)$ is always a $C_0(X)$ -algebra with $(C_0(T, B))_x = C_0(T, B_x)$. One defines $\mathfrak{A}_X B$, which, abusing notation slightly, will be abbreviated to $\mathfrak{A}B$, to be the quotient $C_b(T, B)/C_0(T, B)$; then $\mathfrak{A}B$ is a $C_0(X)$ -algebra. A $C_0(X)$ -asymptotic morphism is defined to be a $C_0(X)$ -morphism $\phi : A \rightarrow \mathfrak{A}B$.

Now suppose that A, B are Γ -algebras. We would like $C_b(T, B), \mathfrak{A}B$ to be Γ -algebras in a natural way so that we can define Γ -equivariant asymptotic morphisms from A to $\mathfrak{A}B$. As we will see, there is a technical difficulty in defining the appropriate Γ -actions, and indeed, even in the group case of [12], continuous versions of $C_b(T, B), \mathfrak{A}B$ are required. The C^* -algebras $(C_b(T, B))_x$ make sense, of course,

since $C_b(T, B)$ is now a $C_0(X)$ -algebra. The problem is to obtain a natural Γ -action on $C_b(T, B)$: how does one define the $\alpha_\gamma : (C_b(T, B))_{s(\gamma)} \rightarrow (C_b(T, B))_{r(\gamma)}$? To deal with this it is natural to try to replace $C_b(T, B)_x$ by $C_b(T, B_x)$ and $C_b(T, B)^\#$ by the bundle $C_b(T, B^\#) = \sqcup_{x \in X} C_b(T, B_x)$; for, using the Γ -action on B , $C_b(T, B^\#)$ is a Γ -space of C^* -algebras in the natural way:

$$(4.1) \quad \alpha_\gamma(h_{s(\gamma)})(t) = \alpha_\gamma(h_{s(\gamma)}(t))$$

where, of course, $h_{s(\gamma)} \in C_b(T, B_{s(\gamma)})$. For each x , there is a canonical homomorphism $R_x : C_b(T, B) \rightarrow C_b(T, B_x)$ given by: $R_x(F)(t) = F(t)_x$. Note that $R_x(fF) = f(x)R_xF$ ($f \in C_0(X)$). Since $R_x(I_x C_b(T, B)) = 0$, the map R_x descends to a $*$ -homomorphism, also denoted R_x from $C_b(T, B)_x$ into $C_b(T, B_x)$. Since $R_x(C_0(T, B)) \subset C_0(T, B_x)$, it also induces a $*$ -homomorphism, $i_x : \mathfrak{A}B \rightarrow \mathfrak{A}(B_x)$. Then i_x determines a $*$ -homomorphism, also denoted $i_x : (\mathfrak{A}B)_x \rightarrow \mathfrak{A}(B_x)$: $i_x(\overline{F_x}) = \overline{R_x F}$ where $\overline{F} = F + C_0(T, B)$ and for $g \in C_b(T, B_x)$, we set $\overline{g} = g + C_0(T, B_x)$. If we knew that R_x, i_x were onto isomorphisms, then we could identify $C_b(T, B)_x$ with $C_b(T, B_x)$ and $(\mathfrak{A}B)_x$ with $\mathfrak{A}(B_x)$ and be able to define (at least algebraically) actions of Γ on $C_b(T, B), \mathfrak{A}B$.

Unfortunately, the C^* -algebra $C_b(T, B)$ is too big for this to work (as we will see below). However, there is a very useful, simple relation between sections of the bundles $C_b(T, B)^\#$ and $C_b(T, B^\#)$ which we now describe. For each $F \in C_b(T, B) = C_0(X, C_b(T, B)^\#)$ (Theorem 1), define a section RF of the bundle $C_b(T, B^\#)$ by setting: $RF(x) = R_x F = R_x F_x$. Let $S_0(X, C_b(T, B^\#))$ be the C^* -algebra of sections of the bundle $C_b(T, B^\#)$ that vanish at infinity. The support $\text{supp } H$ of a section $H \in S_0(X, C_b(T, B^\#))$ is the closure in X of the set $\{x \in X : H(x) \neq 0\}$. For $F \in C_b(T, B)$, the support $\text{supp } F$ of F is the $(X-)$ support of the section $x \rightarrow F_x$ (not the support of F as a function of t).

Proposition 3. *The map $R : C_b(T, B) \rightarrow S_0(X, C_b(T, B^\#))$ is a support preserving $*$ -isomorphism.*

Proof. The only non-trivial part of the proof that R is a $*$ -isomorphism is to show R is one-to-one. Suppose then that $RF = 0$. Then for fixed t , $F(t)_x = 0$ in B_x for all x . So $F(t) = 0$ by Theorem 1, and so $F = 0$. Now let $F \in C_b(T, B)$ be general. Since R_x is norm-decreasing, we obtain that $\text{supp } RF \subset \text{supp } F$. Now suppose that $\text{supp } RF \neq \text{supp } F$. Then there exists an open set V in X such that $V \cap \text{supp } RF = \emptyset$ and $f \in C_c(V)$ such that $fF \neq 0$. Then $R(fF) = f(RF) = 0$ and we contradict the $*$ -isomorphism property. \square

In most of what follows, we will replace $C_b(T, B)$ by a smaller $C_0(X)$ -subalgebra \mathfrak{B} that contains $C_0(T, B)$. The constructions above for $C_b(T, B)$ go through for \mathfrak{B} . Let $\mathfrak{B}_T = \sqcup_{x \in X} R_x \mathfrak{B}$, a bundle of C^* -algebras that is a subbundle of $C_b(T, B^\#)$. As above, we obtain, for each x , a $*$ -homomorphism $R_x : \mathfrak{B} \rightarrow R_x \mathfrak{B}$, which descends to $R_x : \mathfrak{B}_x \rightarrow R_x \mathfrak{B}$. Then $\mathfrak{B}/C_0(T, B)$ is a $C_0(X)$ -algebra, with $(\mathfrak{B}/C_0(T, B))_x = \mathfrak{B}_x/C_0(T, B_x)$. We obtain a canonical homomorphism $i_x : (\mathfrak{B}/C_0(T, B))_x \rightarrow R_x(\mathfrak{B}_x)/C_0(T, B_x)$. We write $\mathfrak{A}_{\mathfrak{B}}B, \mathfrak{A}_{\mathfrak{B}}B_x$ in place of $\mathfrak{B}/C_0(T, B), R_x(\mathfrak{B}_x)/C_0(T, B_x)$. Note that R_x, i_x are onto, but unfortunately, are not usually injective.

For example, consider the case where $X = [0, 1]$ and $B = C([0, 1])$. Then the function F on T , where $F(t)(x) = \sin(tx)$, belongs to $C_b(T, B)$ using the mean-value theorem, and $R_0(F)(t) = \sin(t0) = 0$. So $R_0 F = 0$. But $F_0 \neq 0$. For otherwise, $F = fF'$ for some $f \in I_0, F' \in C_b(T, B)$, and so $\sup_{t \in T} |\sin(tx)| \rightarrow 0$ as $x \rightarrow 0$.

But if x is not zero, then $\sup_{t \in T} |\sin(tx)| = 1$. It is obvious that the image \overline{F}_0 of F in $(\mathfrak{A}B)_0$ is non-zero yet $i_0(\overline{F}_0) = 0$ in $\mathfrak{A}B_0$. So i_0 is not injective as well.

We now look at the question: when are the R_x 's $*$ -isomorphisms for \mathfrak{B} as above? If the latter holds, then every R_x must be an isometry on \mathfrak{B}_x , and it follows that the map $x \rightarrow \|R_x F\|$ is upper semicontinuous for all $F \in \mathfrak{B}$. Here is the converse.

Proposition 4. *Suppose that the map $y \rightarrow \|R_y F\|$ is upper semicontinuous for all $F \in \mathfrak{B}$. Then for every $y \in X$, R_y is a $*$ -isomorphism on \mathfrak{B}_y .*

Proof. Let $F \in \mathfrak{B}$, $y \in X$ and $\epsilon > 0$. Since $\|R_x F\| \leq \|F_x\| \rightarrow 0$ as $x \rightarrow \infty$, there exists a compact subset C of X such that $\|R_x F\| < \epsilon$ for all $x \in X \setminus C$. Suppose that $R_y F = 0$. By upper semicontinuity, there exists an open neighborhood U of y such that $\|R_x F\| < \epsilon$ for all $x \in U$. We can suppose that $U \subset C$. Let $h \in C_c(X)$ be such that $0 \leq h \leq 1$, $h(y) = 0$ and $h(x) = 1$ for all $x \in C \setminus U$. Also $\|R_x(F - hF)\| = 0$ if $x \in C \setminus U$, $\leq \|R_x F\| < \epsilon$ if $x \in U$, $< \epsilon$ if $x \in X \setminus C$. So $\|R_x(F - hF)\| < \epsilon$ for all $x \in X$. Note next that $\|R_x(F - hF)\| = (1 - h(x))\|R_x F\|$. So for each t , $\|F(t)_x - h(x)F(t)_x\| < \epsilon$ for all x , and so by Theorem 1, $\|F(t) - hF(t)\| \leq \epsilon$. So $\|F - hF\| \leq \epsilon$. Since $h \in I_y$, $\|F_y\| \leq \epsilon$, and since ϵ was arbitrary, $F_y = 0$. So $R_y : \mathfrak{B}_y \rightarrow C_b(T, B_y)$ is injective and so isometric. \square

We note that under the condition of Proposition 4, every i_y is also isometric. Now let B be a Γ -algebra. We have the canonical action of Γ on the bundle \mathfrak{B}_T : $\alpha_\gamma(F_{s(\gamma)})(t) = \alpha_\gamma(F(t)_{s(\gamma)})$, $F \in \mathfrak{B}$ provided that \mathfrak{B} is Γ -invariant, i.e. $\alpha_\gamma(R_{s(\gamma)}\mathfrak{B}) = R_{r(\gamma)}\mathfrak{B}$ for all γ . In that situation, we look for an action β of Γ on \mathfrak{B} - not necessarily continuous - related to α on \mathfrak{B}_T ; precisely, we want β to *cover* α in the sense that for each γ , the following diagram commutes:

$$\begin{array}{ccc} \mathfrak{B}_{s(\gamma)} & \xrightarrow{\beta_\gamma} & \mathfrak{B}_{r(\gamma)} \\ R_{s(\gamma)} \downarrow & & \downarrow R_{r(\gamma)} \\ R_{s(\gamma)}\mathfrak{B}_{s(\gamma)} & \xrightarrow{\alpha_\gamma} & R_{r(\gamma)}\mathfrak{B}_{r(\gamma)} \end{array}$$

i.e.,

$$(4.2) \quad R_{r(\gamma)}\beta_\gamma = \alpha_\gamma R_{s(\gamma)}$$

on $\mathfrak{B}_{s(\gamma)}$. If this condition is satisfied and if β restricted to $C_0(T, B)$ is the canonical Γ -algebra action on $C_0(T, B)$ ((2.4)), then we say that \mathfrak{B} has a *covering* Γ -action. In the case of a continuous covering Γ -action, we can extend the continuity condition for a continuous functor \mathbf{F} from finite intervals $I = [a, b]$ to the infinite interval T . For such an I , the canonical Γ -action on IB (also as in (2.4)) will be denoted by $\gamma \rightarrow \alpha'_\gamma$. It is determined by: $R_{r(\gamma)}\alpha'_\gamma = \alpha_\gamma R_{s(\gamma)}$, where $\alpha'_\gamma(h_{s(\gamma)})(t) = \alpha_\gamma(h(t)_{s(\gamma)})$ (using R_x to identify $(IB)_x$ with IB_x).

Proposition 5. *Let $\mathfrak{B} \subset C_b(T, B)$ have a covering Γ -action and \mathbf{F} be a continuous functor as in the third section. Then for all $k \in \mathbf{F}(\mathfrak{B})$, the function $\hat{k} \in C_b(T, \mathbf{F}(\mathbf{B}))$, and if $\mathfrak{B} = C_0(T, B)$, then $\hat{k} \in C_0(T, \mathbf{F}(\mathbf{B}))$.*

Proof. Let I be a closed bounded interval inside T and $\rho : \mathfrak{B} \rightarrow IB$ be the restriction map. We show first that ρ is a Γ -homomorphism. Note first that ρ is a $C_0(X)$ -homomorphism: for since $fF(t) = f \cdot F(t)$ ($f \in C_0(X)$, $F \in \mathfrak{B}$) for all t ,

we get $\rho(fF) = f\rho(F)$. Also, for $t \in I$, $R_x \rho_x F_x(t) = F(t)_x = R_x F_x(t)$. Using the above and (4.2), for $t \in I$,

$$\begin{aligned} R_{r(\gamma)} \rho_{r(\gamma)} \beta_\gamma F_{s(\gamma)}(t) &= R_{r(\gamma)} \beta_\gamma F_{s(\gamma)}(t) = \alpha_\gamma [R_{s(\gamma)} F_{s(\gamma)}(t)] \\ &= \alpha_\gamma [R_{s(\gamma)} \rho_{s(\gamma)} F_{s(\gamma)}(t)] = R_{r(\gamma)} \alpha'_\gamma \rho_{s(\gamma)} F_{s(\gamma)}(t). \end{aligned}$$

Since $R_{r(\gamma)}$ is a bijection, we obtain $\rho_{r(\gamma)} \beta_\gamma = \alpha'_\gamma \rho_{s(\gamma)}$, so that ρ is a Γ -homomorphism. With I a single point t_0 , we get that ev_{t_0} is also a Γ -homomorphism. Now let $I = [a, b] \subset T$ and ρ be as above. For $t \in I$, $ev_t = ev_t \circ \rho$, and so for $k \in \mathbf{F}(\mathfrak{B})$, $\hat{k}(t) = \mathbf{F}(ev_t)(k) = \mathbf{F}(ev_t)(\mathbf{F}(\rho)(k))$ which is continuous in t by the continuity of \mathbf{F} . So $\hat{k} \in C_b(T, \mathbf{F}(B))$. The last part is just (3.11). \square

One natural way in which a covering Γ -action on some \mathfrak{B} can arise is when, for each x , R_x is an isomorphism on \mathfrak{B}_x and \mathfrak{B}_T is Γ -invariant, in which case we can identify the bundles \mathfrak{B}^\sharp and \mathfrak{B}_T and obtain a Γ -action on \mathfrak{B} : effectively $\beta = \alpha$ in this case. An example of this is the situation of Proposition 6 below. However, there are many cases where the R_x 's are not isomorphisms but we can still find a covering action on some reasonable \mathfrak{B} . For instance, in the example above with $X = [0, 1]$, where we take $\Gamma = X$ (a groupoid of units), we can take $\mathfrak{B} = C[0, 1]$ and $\beta_x(F_x) = F_x!$. Of course, this example is trivial, but as we will see in Theorem 5, such a \mathfrak{B} always exists under very general circumstances.

Suppose now that Γ has local $r - G$ -sets, and that \mathfrak{B} has a covering Γ -action. By Proposition 2, \mathfrak{B}^{cont} is a $\Gamma - C^*$ -algebra. By definition, the β action restricts to give the canonical Γ -action on $C_0(T, B)$ which is continuous. So $C_0(T, B) \subset \mathfrak{B}^{cont}$. So we can define $\mathfrak{A}_{\mathfrak{B}^{cont}} B = \mathfrak{B}^{cont} / C_0(T, B)$. If A is also a $\Gamma - C^*$ -algebra, then an *equivariant asymptotic morphism* from A to B (relative to \mathfrak{B}^{cont}) is just a Γ -homomorphism $\phi : A \rightarrow \mathfrak{A}_{\mathfrak{B}^{cont}} B$. In the locally compact group case, one takes $\mathfrak{B} = C_b(T, B)$ and the asymptotic algebra $\mathfrak{A}_{\mathfrak{B}^{cont}} B = C_b(T, B)^{cont} / C_0(T, B)$ is the same as the $\mathfrak{A}B$ of [12, p.7]. In that case, there is a descent functor for $\Gamma - C^*$ -algebras using as morphisms homotopy classes of Γ -homomorphisms into the asymptotic algebra ([12, Theorem 4.12]). However, since, for completely general Γ , we do not have available a canonical \mathfrak{B}^{cont} , it does not make sense to talk of “the functor $B \rightarrow \mathfrak{A}_{\mathfrak{B}^{cont}} B$ ”. (However, for a very wide class of groupoids Γ , we do obtain a canonical asymptotic algebra and a functor from Theorem 5 below - it would be interesting to know if the complete theory of the descent functor can be developed for this class of groupoids as in [12].) Instead at present, we avoid a functorial description of the descent functor and give a direct, weaker version of the descent homomorphism which is adequate for a number of applications.

Theorem 4. *Suppose that Γ has local $r - G$ -sets, and that \mathfrak{B} is a $C_0(X)$ subalgebra of $C_b(T, B)$ containing $C_0(T, B)$ and with a covering Γ -action. Let A be a $\Gamma - C^*$ -algebra and $\phi : A \rightarrow \mathfrak{A}_{\mathfrak{B}^{cont}} B$ be a Γ -homomorphism. Then there exists a canonical descent homomorphism (dependent on \mathfrak{B}) given by: $i \circ \mathbf{F}(\phi) : C^*(\Gamma, A) \rightarrow \mathfrak{A}C^*(\Gamma, B)$.*

Proof. The proof is effectively the same as for the group case ([12, Theorem 4.12]). Let \mathbf{F} be as in Theorem 3. From the exactness of \mathbf{F} , we get the short exact sequence:

$$0 \rightarrow \mathbf{F}(C_0(T, B)) \rightarrow \mathbf{F}(\mathfrak{B}^{cont}) \rightarrow \mathbf{F}(\mathfrak{A}_{\mathfrak{B}^{cont}} B) \rightarrow 0.$$

We also have the short exact sequence for the ordinary C^* -algebra $\mathbf{F}(B)$:

$$0 \rightarrow C_0(T, \mathbf{F}(B)) \rightarrow C_b(T, \mathbf{F}(B)) \rightarrow \mathfrak{A}\mathbf{F}(B) \rightarrow 0.$$

Continuity (Proposition 5) gives $*$ -homomorphisms $i_0 : \mathbf{F}(C_0(T, B)) \rightarrow C_0(T, \mathbf{F}(B))$, $i_b : \mathbf{F}(\mathfrak{B}^{cont}) \rightarrow C_b(T, \mathbf{F}(B))$ with i_0 the restriction of i_b to $\mathbf{F}(C_0(T, B))$, and these induce a $*$ -homomorphism $i : \mathbf{F}(\mathfrak{A}_{\mathfrak{B}^{cont}} B) \rightarrow \mathfrak{A}\mathbf{F}(B)$. Next we have a $*$ -homomorphism $\mathbf{F}(\phi) : \mathbf{F}(A) \rightarrow \mathbf{F}(\mathfrak{A}_{\mathfrak{B}^{cont}} B)$. So $i \circ \mathbf{F}(\phi) : \mathbf{F}(A) \rightarrow \mathfrak{A}\mathbf{F}(B)$ is a $*$ -homomorphism. \square

Before discussing our main theorem giving a canonical \mathfrak{B} we look at a situation in which there is a very simple \mathfrak{B} available. For motivation, Proposition 4 suggests that we should look for a \mathfrak{B} with elements F for which the map $x \rightarrow \|R_x F\|$ is upper semicontinuous. I have been unable to find a canonical such \mathfrak{B} in general. However, in cases that arise in practice - in particular, when Γ is discrete, or when Γ is a locally compact group (the case of [12]) or when $B = \mathcal{C}(E) = C_0(\mathbb{R}) \hat{\otimes} C_0(E, \text{Cliff}(E))$ (E a Γ -vector bundle) - there is a natural such \mathfrak{B} available. (The last of these cases is needed for the groupoid version of the infinite dimensional Bott periodicity theorem of Higson, Kasparov and Trout ([14]).) Intuitively, we wish to exclude from \mathfrak{B} functions such as $\sin(xt)$ by requiring *uniformly continuity* in the X -direction. This requires a strong condition on B , but the groupoid Γ has to satisfy only the weak local $r - G$ -set condition of Proposition 2.

We know that B is isomorphic to the C^* -algebra $C_0(X, B^\sharp)$ of the C^* -bundle (B^\sharp, p) . We assume now that this C^* -bundle B^\sharp is a locally trivial C^* -bundle with (C^* -algebra) fiber C . Precisely, a chart (U, η) is given by an open subset U of X together with a fiber preserving homeomorphism η from $p^{-1}(U)$ onto $U \times C$ with each η_x a $*$ -isomorphism from B_x onto C ($x \in U$). Local triviality means that the chart sets U cover X . (In particular, no structure group condition is imposed.) For such a chart (U, η) and $F \in C_b(T, B)$, $R_x F \in C_b(T, B_x)$ and so for $x \in U$, $\eta_x \circ R_x F \in C_b(T, C)$. We say that F is *uniformly continuous* (for X) if the map $\eta R(F) : x \rightarrow \eta_x \circ R_x F$ is continuous from U to $C_b(T, C)$ for every chart (U, η) . It is easily checked that the set $\mathfrak{B} = UC_b(T, B)$ of uniformly continuous functions is a C^* -algebra, and a $C_0(X)$ -subalgebra of $C_b(T, B)$. Note that the equality $R_x \mathfrak{B} = C_b(T, B_x)$ below shows that this \mathfrak{B} is “big”.

Proposition 6. *Let $\mathfrak{B} = UC_b(T, B)$. Then $C_0(T, B) \subset \mathfrak{B}$, and for $F \in \mathfrak{B}$, the map $x \rightarrow \|R_x F\|$ is continuous. The maps R_x, i_x are $*$ -isomorphisms. Further $R_x(\mathfrak{B}) = C_b(T, B_x)$, and \mathfrak{B} is Γ -invariant (so that trivially \mathfrak{B} has a covering Γ -action).*

Proof. Let $k \in C_0(T, B)$. To show that $k \in UC_b(T, B)$, we can suppose that $k = h \otimes b$ where $h \in C_c(T)$ and $b \in C_c(X)B$. Using a partition of unity, we can suppose that there is a chart (U, η) and a compact subset K of U such that $k_x = 0$ for all $x \in X \setminus K$. Then $\eta R(k) \in C_c(T \times U, C)$, and $k \in \mathfrak{B}$. So $C_0(T, B) \subset \mathfrak{B}$. Now let $F \in \mathfrak{B}$. Then on U , the map $x \rightarrow \|\eta_x \circ R_x F\| = \|R_x F\|$ is continuous. By Proposition 4, each R_x, i_x is an isomorphism. By definition, $R_x(\mathfrak{B}) \subset C_b(T, B_x)$. To show equality, we just have to show that if $H \in C_c(U, C_b(T, C))$ then there exists $F \in C_b(T, B)$ such that $\eta R(F) = H$ (so that $F \in \mathfrak{B}$). For then we can take any $\phi \in C_b(T, B_x)$, take $g \in C_c(U)$ with $g(x) = 1$ and $H(y) = g(y)\eta_x \circ \phi$ to get $R_x F = \phi$.

To show that such an F exists, fix t . The map $y \rightarrow H_y(t) \in C$ is continuous on U , and so $y \rightarrow \eta_y^{-1} H_y(t)$ is a continuous section of B^\sharp supported on U . By Theorem 1, there exists $b_t \in B$ such that $(b_t)_y = \eta_y^{-1} H_y(t)$. Define $F(t) = b_t$. Then F is bounded since $\|(b_t)_y\| \leq \|H_y(t)\| \leq \|H\|_\infty$. Last F is continuous. Indeed, given

$\epsilon > 0$, there exists a $\delta > 0$ such that for all $y \in U$, $\|H_y(t) - H_y(s)\| < \epsilon$ whenever $|t - s| < \delta$. Let $|t - s| < \delta$. Then for all y , $\|(b_t)_y - (b_s)_y\| < \epsilon$. Now take the supremum over y to get F continuous at t . \square

It follows from Theorem 4 that if Γ has local r - G -sets, then $UC_b(T, B)$ determines a descent homomorphism: this can be regarded as the natural descent homomorphism for such a special B . We now show that under a very mild condition on Γ and with B completely general, there *always* exists a canonical \mathfrak{B} with a covering Γ -action giving a descent homomorphism. The algebra \mathfrak{B} is functorial. The condition that we need on Γ is that it have *local G -sets*, i.e. local transversals for r and s simultaneously. The algebra \mathfrak{B} consists (roughly) of those functions with a transversally continuous action, and going to \mathfrak{B}^{cont} then gives continuity of the action in every direction. We now make these ideas precise.

A subset W of Γ is called a G -set (cf. [32, p.10]) if the restrictions r_W, s_W of r, s to W are homeomorphisms onto open subsets of X . The family of G -sets in Γ is denoted by \mathcal{G} .

Proposition 7. *\mathcal{G} is an inverse semigroup under pointwise product and inversion.*

Proof. The discrete case is given in [25, Proposition 2.2.3]. For the topological conditions, we need to show that for $W, W_1, W_2 \in \mathcal{G}$, the bijections $r_{W_1 W_2}, s_{W_1 W_2}, r_{W^{-1}}, s_{W^{-1}}$ are homeomorphism onto open subsets of X . We will prove this for $r_{W_1 W_2}$ leaving the rest to the reader. First, $r(W_1 W_2) = r_{W_1} s_{W_1}^{-1}(r(W_2) \cap s(W_1))$ which is open in X . Next let $x_n \rightarrow x$ in $r(W_1 W_2)$. We write uniquely $x_n = r(w_1^n w_2^n), x = r(w_1 w_2)$ where $w_1^n, w_1 \in W_1, w_2^n, w_2 \in W_2$. Then $(r_{W_1})^{-1}(x_n) = w_1^n \rightarrow (r_{W_1})^{-1}(x) = w_1$, and similarly, $w_2^n \rightarrow w_2$. So $w_1^n w_2^n \rightarrow w_1 w_2$ and $r_{W_1 W_2}^{-1}$ is continuous. \square

We note that if $W \in \mathcal{G}$, the map $r_W s_W^{-1}$ is a homeomorphism from $s(W)$ onto $r(W)$, and there is a nice formula for $r_W s_W^{-1}$:

$$r_W s_W^{-1}(x) = WxW^{-1}.$$

We will say that Γ has *local G -sets* if $\cup \mathcal{G} = \Gamma$, i.e. every element of Γ belongs to a G -set. This property is satisfied by most groupoids that arise in practice (e.g. r -discrete groupoids, transformation group groupoids, many (all?) Lie groupoids). For motivation for the following, suppose that $W \in \mathcal{G}$. Suppose that $\mathfrak{B} \subset C_b(T, B)$ has a covering Γ -action β that makes it into a Γ -algebra. For $F \in \mathfrak{B}$, the map $r(\gamma) \rightarrow \beta_\gamma(F_{s(\gamma)})$ is continuous and so will come from an element F' of $C_b(T, B)$ (at least after multiplying by a function in $C_c(s(W))$). Now from Proposition 3, we can recover F' from RF' and at the R -level, we do have the action α_γ . The idea then is to consider functions F for which there is an F' that goes down under R to the function $r(\gamma) \rightarrow \alpha_\gamma(R_{s(\gamma)} F)$ and define $\beta_\gamma(F_{s(\gamma)}) = F'_{r(\gamma)}$. We also insist that this definition is independent of the choice of W . This does not ensure a continuous action but only one continuous along the G -sets. However, the algebra of such functions is the largest subalgebra $C_b^\Gamma(T, B)$ of $C_b(T, B)$ admitting a reasonable covering action that is continuous along the G -sets. We now develop the theory of $C_b^\Gamma(T, B)$.

We identify $C_b(T, B)$ with $C_0(X, (C_b(T, B))^\#)$. Abbreviate $C_b(T, B)$ to C_b , and let $F \in C_b$. We say that $F \in C_c^G = C_c^G(\Gamma, B)$ if:

- (1) $F \in C_c(X)C_b$;

- (2) for all $W \in \mathcal{G}$ with $\text{supp } F \subset s(W)$, there exists $F^W \in C_b$ such that

$$RF^W = R_W F$$

where $R_W F(x) = \alpha_\gamma R_{s(\gamma)} F$ if $x = r(\gamma)$ for some $\gamma \in W$, and is 0 otherwise;

- (3) (uniqueness) if $W_1, W_2 \in \mathcal{G}$, $\gamma_0 \in W_1 \cap W_2$ and $\text{supp } F \subset s(W_1) \cap s(W_2)$, then

$$(F^{W_1})_{r(\gamma_0)} = (F^{W_2})_{r(\gamma_0)}.$$

We say that $F \in C_b^\Gamma = C_b^\Gamma(T, B)$ if $fF \in C_c^G$ for all $f \in C_c(X)$.

Theorem 5. $C_b^\Gamma(T, B)$ is a $C_0(X)$ -subalgebra of $C_b(T, B)$ with a covering Γ -action given by:

$$(4.3) \quad \beta_{\gamma_0} F_{s(\gamma_0)} = (fF)_{r(\gamma_0)}^W$$

where $W \in \mathcal{G}$, $\gamma_0 \in W$ and $f \in C_c(s(W))$ is such that $f(s(\gamma_0)) = 1$.

Proof. We prove the theorem in five stages.

(a) Let $F \in C_c^G$ and $W \in \mathcal{G}$ with $\text{supp } F \subset s(W)$. Then F^W is unique, and $\text{supp } F^W = W(\text{supp } F)W^{-1}$. Also $\|F^W\| = \|F\|$.

The uniqueness of F^W follows from Proposition 3. The same proposition gives that $\text{supp } F = \text{supp } RF$, $\text{supp } R_W F = \text{supp } F^W$. Next, since α_γ is an isometry, for $\gamma \in W$, $R_{s(\gamma)} F \neq 0$ if and only if $R_W F(r(\gamma)) \neq 0$, so that $R_x F \neq 0$ if and only if $R_W F(WxW^{-1}) \neq 0$ for $x \in s(W)$. Closing up gives $\text{supp } R_W F = W(\text{supp } RF)W^{-1}$. For the last part, use Proposition 3 and the fact that the α_γ 's are *-isomorphisms.

(b) Let $F \in C_c^G$ and $W \in \mathcal{G}$ with $\text{supp } F \subset s(W)$. Then $F^W \in C_c^G$, and for $V \in \mathcal{G}$ with $\text{supp } F^W \subset s(V)$, $(F^W)^V = F^{VW}$. Further, $F^* \in C_c^G$ and $(F^*)^W = (F^W)^*$.

By (a), $\text{supp } F^W = W(\text{supp } F)W^{-1} \subset V^{-1}V$, and so $\text{supp } F \subset s(VW)$. Now $R_V F^W(x) = \alpha_{\gamma_1} R_{s(\gamma_1)} F^W$ if $x = r(\gamma_1)$ for some $\gamma_1 \in V$, and is 0 otherwise. Next $\alpha_{\gamma_1} R_{s(\gamma_1)} F^W = \alpha_{\gamma_1} (\alpha_{\gamma_2} R_{s(\gamma_2)} F)$ if $s(\gamma_1) = r(\gamma_2)$ for some (unique) $\gamma_2 \in W$ and is 0 otherwise. Since $\alpha_{\gamma_1} (\alpha_{\gamma_2} R_{s(\gamma_2)} F) = \alpha_{\gamma_1 \gamma_2} R_{s(\gamma_1 \gamma_2)} F$ and $r(\gamma_1) = r(\gamma_1 \gamma_2)$, it follows that $R_V F^W = RF^{VW}$. So $(F^W)^V = F^{VW}$. The uniqueness condition (3) of the definition of C_c^G with respect to F^W follows from the corresponding property for F . The last part of the proof of (b) is easy.

(c) Let $F \in C_b^\Gamma$, $W \in \mathcal{G}$, $f \in C_c(s(W))$ and $\text{supp } F \subset s(W)$ be compact. Then $(fF)^W = f^W F^W$ and belongs to C_b^Γ , and $R_W(fF) = R(f^W F^W)$ where $f^W \in C_c(r(W))$ is given by: $f^W(y) = f(W^{-1}yW)$.

Note that by definition, $fF \in C_b^\Gamma$. Next, if $\gamma \in W$ and $y = r(\gamma)$, then $R_W(fF)(y) = \alpha_\gamma(R_{s(\gamma)}(fF)) = f(s(\gamma))R_W(F)(y) = f^W(r(\gamma))R_W F(r(\gamma)) = R_y(f^W F^W)$. Both $R_W(fF)$, $R(f^W F^W)$ vanish at y if $y \notin r(W)$, and so $R_W(fF) = R(f^W F^W)$. By (a), $(fF)^W = f^W F^W$. Next we show that $(fF)^W \in C_b^\Gamma$. For let $g \in C_c(X)$ and $h \in C_c(r(W))$ be such that $h = 1$ on $\text{supp } F^W = W(\text{supp } F)W^{-1}$, a compact subset of $r(W)$ (using (a)). Then $g(fF)^W = (gh)(fF)^W = ((gh)^{W^{-1}})^W (fF)^W = ((gh)^{W^{-1}} fF)^W \in C_c^G$ by (b). So $(fF)^W \in C_b^\Gamma$.

(d) C_b^Γ is a $C_0(X)$ -subalgebra of C_b that contains $C_0(T, B)$.

Trivially, $0 \in C_b^\Gamma$. Let $F_1, F_2 \in C_b^\Gamma$, $f \in C_c(X)$, $W \in \mathcal{G}$ with $C = \text{supp}(f(F_1 + F_2)) \subset s(W)$. Choose $f' \in C_c(s(W))$ so that $f' = 1$ on C . Then $f(F_1 + F_2) =$

$f'fF_1 + f'fF_2$ with $\text{supp } f'fF_1 \cup \text{supp } f'fF_2 \subset s(W)$. Since $F_1, F_2 \in C_b^\Gamma$, we get $f'fF_1, f'fF_2 \in C_c^G$. Then $R_W(fF_1 + fF_2) = R_W(f'fF_1) + R_W(f'fF_2) = R((f'fF_1)^W + (f'fF_2)^W)$ and we obtain $(fF_1 + fF_2)^W = (f'fF_1)^W + (f'fF_2)^W$. For uniqueness, let $W' \in \mathcal{G}$, $C \subset s(W')$ and $\gamma_0 \in W \cap W'$. Then we can choose $f' \in C_c(s(W) \cap s(W'))$ and obtain uniqueness using $((f'fF_1)^W)_{r(\gamma_0)} = ((f'fF_1)^{W'})_{r(\gamma_0)}, ((f'fF_2)^W)_{r(\gamma_0)} = ((f'fF_2)^{W'})_{r(\gamma_0)}$. So $F_1 + F_2 \in C_b^\Gamma$. Next, $F_1F_2 \in C_b^\Gamma$. The proof is similar to that for the sum above. Let $W \in \mathcal{G}$ with $C = \text{supp } fF_1F_2 \subset s(W)$. Choose $f' \in C_c(s(W))$ with $f' = 1$ on C . Then $fF_1F_2 = (f'fF_1)(f'F_2)$ and we can use $R_W(fF_1F_2) = R_W(f'fF_1)R_W(f'F_2)$ to get $(fF_1F_2)^W = (f'fF_1)^W(f'F_2)^W$. The remaining details are left to the reader, as is also the proof that $F_1^* \in C_b^\Gamma$ (use (b)). So C_b^Γ is a $*$ -subalgebra of C_b .

Next we show that C_b^Γ is complete. Let $F_n \rightarrow F$ in C_b with every $F_n \in C_b^\Gamma$. We show that $F \in C_b^\Gamma$. Let $f \in C_c(X)$ and $W \in \mathcal{G}$ be such that $\text{supp } fF \subset s(W)$. We can suppose that $f \in C_c(s(W))$. Let $g \in C_c(s(W))$ be 1 on the support of f . Using (a) and Proposition 3, $\|gF_n - gF_m\| = \|R(gF_n - gF_m)\| = \|(gF_n)^W - (gF_m)^W\| \rightarrow 0$ as $n, m \rightarrow \infty$. So there exists $F'' \in C_b$ such that $(gF_n)^W \rightarrow F''$ in norm. By the continuity of R_W, R , we get for $\gamma \in W$, $R_W(fF)(r(\gamma)) = \lim R_W(f(gF_n))(r(\gamma)) = \lim R_{r(\gamma)}(f^W(gF_n)^W) = R_{r(\gamma)}(f^WF'')$. We take $(fF)^W = f^WF''$. Uniqueness is proved using uniqueness for the F_n 's and a simple limit argument. To prove that $C_0(T, B) \subset C_b^\Gamma$, let $F \in C_0(T, B)$, $f \in C_c(X)$, $W \in \mathcal{G}$, $\text{supp}(fF) \subset s(W)$. Then we take $(fF)^W(r(\gamma)) = \alpha_\gamma(fF)_{s(\gamma)}$ ($\gamma \in W$), where α is the canonical Γ -algebra action. Uniqueness is obvious. So $fF \in C_c^G$ and $F \in C_b^\Gamma$.

Last we have to show that C_b^Γ is a $C_0(X)$ -algebra. Let $F \in C_b^\Gamma$ and $h \in C_0(X)$. Trivially, if $f \in C_c(X)$ then $f(hF) = (fh)F \in C_c^G$. So $hF \in C_b^\Gamma$. To prove that the action of $C_0(X)$ on C_b^Γ is non-degenerate use the fact that $C_b = C_0(X)C_b$.

(e) The β_γ 's give a covering Γ -action on C_b^Γ .

We first show that, for given $F \in C_b^\Gamma$, the right hand side of (4.3) is well-defined. Let f, W be as in (4.3) and let $W' \in \mathcal{G}$, $\gamma_0 \in W'$ and $f' \in C_c(s(W'))$ be such that $f'(s(\gamma_0)) = 1$. Suppose first that $W = W'$. Let $g \in C_c(s(W))$ be such that $g = 1$ on $\text{supp } f \cup \text{supp } f'$. Then $fF = fgF$, $f'F = f'gF$, and using (c), $(fF)_{r(\gamma_0)}^W = f^W(r(\gamma_0))(gF)_{r(\gamma_0)}^W = (gF)_{r(\gamma_0)}^W = (f'F)_{r(\gamma_0)}^W$. For the case $W \neq W'$, find $h \in C_c(s(W) \cap s(W'))$ such that $h(s(\gamma_0)) = 1$. Then using uniqueness, $(fF)_{r(\gamma_0)}^W = (hF)_{r(\gamma_0)}^W = (hF)_{r(\gamma_0)}^{W'} = (f'F)_{r(\gamma_0)}^{W'}$. Next we have to show that the right hand side of (4.3) depends only on the coset $F_{s(\gamma_0)}$. So let $F' = F + gF_1$ where $g \in I_{s(\gamma_0)}$ and $F_1 \in C_b^\Gamma$. Then $(fF')_{r(\gamma_0)}^W = (fF)_{r(\gamma_0)}^W + g^W(r(\gamma_0))(fF_1)_{r(\gamma_0)} = (fF)_{r(\gamma_0)}^W$. We now show that $\gamma \rightarrow \beta_\gamma$ defines an algebraic action on C_b^Γ . It is simple, using (b) and the proof of (d) to show that each β_γ is a $*$ -homomorphism.

Next we show that $\beta_{\gamma_0}\beta_{\gamma_1} = \beta_{\gamma_0\gamma_1}$ whenever $s(\gamma_0) = r(\gamma_1)$. Let $F \in C_b^\Gamma$ and let $W \in \mathcal{G}$ contain γ_1 and $f \in C_c(s(W))$ be such that $f(s(\gamma_1)) = 1$. Then $\beta_{\gamma_1}F_{s(\gamma_1)} = (fF)_{r(\gamma_1)}^W$. By (c), $(fF)^W \in C_b^\Gamma$ so that $(fF)_{r(\gamma_1)}^W \in (C_b^\Gamma)_{r(\gamma_1)}$. Now let $V \in \mathcal{G}$ be such that $\gamma_0 \in V$. Since $s(\gamma_0) = r(\gamma_1)$ belongs to $s(V) \cap r(W)$, we can find $g \in C_c(s(V) \cap r(W))$ such that $g(s(\gamma_0)) = 1$. Then $\beta_{\gamma_0}(\beta_{\gamma_1}F_{s(\gamma_1)}) =$

$(g(fF)^W)_{r(\gamma_0)}^V = (((g^{W^{-1}}fF)^W)^V)_{r(\gamma_0)} = ((g^{W^{-1}}fF)^{VW})_{r(\gamma_0\gamma_1)}$ using (b). Noting that $(g^{W^{-1}}f)(s(\gamma_0\gamma_1)) = g(Ws(\gamma_0\gamma_1)W^{-1})f(s(\gamma_1)) = g(r(\gamma_1))f(s(\gamma_1)) = 1$ and $\gamma_0\gamma_1 \in VW$, we see that $((g^{W^{-1}}fF)^{VW})_{r(\gamma_0\gamma_1)}$ is just $\beta_{\gamma_0\gamma_1}F_{s(\gamma_1)}$ and $\beta_{\gamma_0}\beta_{\gamma_1} = \beta_{\gamma_0\gamma_1}$. If $x \in X$, then trivially (using $W = X$), $\beta_x : (C_b^\Gamma)_x \rightarrow (C_b^\Gamma)_x$ is the identity, and it follows that $\gamma \rightarrow \beta_\gamma$ is an algebraic Γ -action on C_b^Γ . To show that the Γ -action is covering, in an obvious notation, $R_{r(\gamma)}(\beta_\gamma F_{s(\gamma)}) = R_{r(\gamma)}(fF)^W = R_W(fF)(r(\gamma)) = \alpha_\gamma(R_{s(\gamma)}(fF)) = \alpha_\gamma R_{s(\gamma)}(fF)_{s(\gamma)} = \alpha_\gamma R_{s(\gamma)}F_{s(\gamma)}$. It is left to the reader to check that β_γ restricted to $C_0(T, B)$ gives the canonical Γ -action on $C_0(T, B)$. \square

Since Γ has local G -sets, it satisfies the condition of Proposition 2, and so $C_b^\Gamma(T, B)^{cont}$ is a Γ -algebra. This can be regarded as the canonical Γ -algebra \mathfrak{B} for groupoids with local G -sets. Indeed, the map $B \rightarrow C_b^\Gamma(T, B)^{cont}$ is functorial in the category of Γ -algebras with Γ -homomorphisms as morphisms. To see this, let B_1, B_2 be Γ -algebras and $\phi : B_1 \rightarrow B_2$ be a Γ -homomorphism. Let $\mathfrak{B}_1 = C_b^\Gamma(T, B_1), \mathfrak{B}_2 = C_b^\Gamma(T, B_2)$. Define $\tilde{\phi} : C_b(T, B_1) \rightarrow C_b(T, B_2)$ by: $\tilde{\phi}(F)(t) = \phi(F(t))$. Then $\psi = \tilde{\phi}$ is a $C_0(X)$ -homomorphism. One readily checks that $R_x(\psi(F)) = \phi_x \circ R_x F$. Let $F \in \mathfrak{B}_1$. Let $W \in \mathcal{G}$, $f \in C_c(X)$ and $\text{supp}(fF) \subset s(W)$. Then for $\gamma \in W$, $\alpha_\gamma R_{s(\gamma)}(f\psi(F)) = \alpha_\gamma R_{s(\gamma)}(\psi(fF)) = \alpha_\gamma(\phi_{s(\gamma)} \circ R_{s(\gamma)}(fF)) = \phi_{r(\gamma)} \circ (\alpha_\gamma R_{s(\gamma)}(fF)) = \phi_{r(\gamma)} \circ R_{r(\gamma)}((fF)^W) = R_{r(\gamma)}(\psi((fF)^W))$. So $(f\psi(F))^W = \psi((fF)^W)$ exists, and uniqueness is easily checked from that for F . So $f\psi(F) \in C_c^\Gamma(T, B_2)$ and $\psi(F) \in C_b^\Gamma(T, B_2)$. Next, if $\gamma_0 \in s(W)$ and $f \in C_c(s(W))$ with $f(s(\gamma_0)) = 1$, we get $\beta_{\gamma_0}(\psi_{s(\gamma_0)}F_{s(\gamma_0)}) = \beta_{\gamma_0}((\psi(F))_{s(\gamma_0)}) = (f\psi(F))_{r(\gamma_0)}^W = (\psi((fF)^W))_{r(\gamma_0)} = \psi_{r(\gamma_0)}(fF)_{r(\gamma_0)}^W = \psi_{r(\gamma_0)}\beta_{\gamma_0}F_{s(\gamma_0)}$, so that ψ is equivariant from \mathfrak{B}_1 to \mathfrak{B}_2 . Last, using equivariance, $\psi : \mathfrak{B}_1^{cont} \rightarrow \mathfrak{B}_2^{cont}$ and the functorial property follows.

REFERENCES

- [1] B. Blackadar, *K-theory for operator algebras*, 2nd edition, MSRI Publications, Vol. 5, Cambridge University Press, Cambridge, 1998.
- [2] É. Blanchard, *Tensor products of $C(X)$ -algebras over $C(X)$* , Recent advances in operator algebras (Orlans, 1992), *Astisque* No. 232 (1995), 81-92.
- [3] É. Blanchard, *Déformations de C^* -algebras de Hopf*, *Bull. Soc. Math. France* 124(1996), 141-215.
- [4] A. Connes, *Sur la théorie non commutative de l'intégration*, *Lecture Notes in Mathematics*, 725(1979), 19-143.
- [5] A. Connes, *Noncommutative Geometry*, Academic Press, Inc., New York, 1994.
- [6] A. Connes, and N. Higson, *Déformations morphismes asymptotiques et K -théorie bivariante*, *C. R. Acad. Sci. Paris, Série 1*, 311(1990), 101-106.
- [7] J. Dixmier, *Von Neumann Algebras*, translation by F. Jellet of: *Les Algèbres d'Opérateurs dans l'Espaces Hilbertien*, Deuxième Édition, Gauthier-Villars, 1969, North-Holland Publishing Company, New York, 1981.
- [8] J. Dixmier, *C^* -algebras*, North-Holland Publishing Company, Amsterdam, 1977.
- [9] Dupré, M. J. and Gillette, R. M., *Banach bundles, Banach modules and automorphisms of C^* -algebras*, Pitman Publishing Inc., Marshfield, Massachusetts, 1983.
- [10] S. Echterhoff and D. P. Williams, *Crossed products by $C_0(X)$ -actions*, *J. Funct. Anal.* 158(1998), 113-151.
- [11] J. M. G. Fell, *An extension of Mackey's method to Banach *-algebraic bundles*, *Mem. Amer. Math. Soc.* 90(1969).

- [12] E. Guenter, N. Higson, and J. Trout, *Equivariant E-theory for C^* -algebras*, Mem. Amer. Math. Soc, 148, no. 703, 2000.
- [13] N. Higson, and G. G. Kasparov, *E-theory and K-theory for groups which act properly and isometrically on Hilbert space*, Invent. Math. 144 (2001), 23-74.
- [14] Higson, N., Kasparov, G. G., and Trout, J., *A Bott Periodicity Theorem for Infinite Dimensional Euclidean Space*, Advances in Mathematics 135 (1998), 1-40.
- [15] Kasparov, G. G., *An index for invariant elliptic operators, K-theory, and representations of Lie groups*, Soviet Math. Dokl. 27(1983), 105-109.
- [16] Kasparov, G. G., *Equivariant KK-theory and the Novikov conjecture*, Invent. Math. 91(1988), 147-201.
- [17] Le Gall, P.-Y., *Théorie de Kasparov équivariante et groupoïdes*, Doctoral Thesis, University of Paris VII, 1994.
- [18] Le Gall, P.-Y., *Théorie de Kasparov équivariante et groupoïdes*, C. R. Acad. Sci. Paris Sér. I, 324(1997), 695-698.
- [19] Le Gall, P.-Y., *Théorie de Kasparov équivariante et groupoïdes I*, K-Theory 16(1999), 361-390.
- [20] Muhly, P. S., *Coordinates in Operator Algebra*, to appear, CBMS Regional Conference Series in Mathematics, American Mathematical Society, Providence.
- [21] Muhly, P. S., Renault, J. N. and Williams, D. P., *Equivalence and isomorphism for groupoid C^* -algebras*, J. Operator Theory 17(1987), 3-22.
- [22] Murphy, G. J., *C^* -algebras and operator theory*, Academic Press, Inc., New York, 1990.
- [23] Nilsen, M., *C^* -bundles and $C_0(X)$ -algebras*, Indiana Univ. Math. J. 45(1996), 463-477.
- [24] Park, E. and Trout, L., *Representable E-theory for $C_0(X)$ -algebras*, J. Funct. Anal. 177(2000), 178-202.
- [25] Paterson, A. L. T., *Groupoids, inverse semigroups and their operator algebras*, Progress in Mathematics, Vol. 170, Birkhäuser, Boston, 1999.
- [26] Paterson, A. L. T., *The equivariant analytic index for proper groupoid actions*, K-Theory 32(2004), 193-230.
- [27] Phillips, N. C., *Equivariant K-theory for proper actions*, Pitman Research Notes in Mathematics, Vol. 178, John Wiley, New York, 1988.
- [28] Phillips, N. C., *Equivariant K-Theory and freeness of group actions on C^* -algebras*, Lecture Notes in Mathematics, vol. 1274, Springer-Verlag, New York, 1987.
- [29] R. Popescu, *E-théorie équivariante et groupoïdes*, C. R. Acad. Sci. Paris Sér I Math 331(2000), 223-228.
- [30] R. Popescu, *Equivariant E-theory for groupoids acting on C^* -algebras*, J. Funct. Anal. 209(2004), 247-292.
- [31] Raeburn, I. and Williams, D. P., *Morita Equivalence and Continuous Trace C^* -algebras*, Mathematical Surveys and Monographs, Vol. 60, American Mathematical Society, Providence, R. I.
- [32] Renault, J. N., *A groupoid approach to C^* -algebras*, Lecture Notes in Mathematics, Vol. 793, Springer-Verlag, New York, 1980.
- [33] Renault, J. N., *Représentation des produits croisés d'algèbres de groupoïdes*, J. Operator Theory 18(1987), 67-97.
- [34] Ramsay, A., *Virtual groups and group actions*, Adv. in Math. 6(1971), 253-322.
- [35] Ramsay, A., *Topologies for measured groupoids*, J. Funct. Anal., 47(1982), 314-343.
- [36] Tu, J. L., *La conjecture de Novikov pour les feuilletages hyperboliques*, K-Theory 16(1999), 129-184.
- [37] Tu, J. L., *La conjecture de Baum-Connes pour les feuilletages moyennables*, K-Theory 17(1999), 215-264.
- [38] Tu, J. L., *The Baum-Connes conjecture and discrete group actions on trees*, K-Theory 17(1999), 303-318.
- [39] Tu, J. L., *The Baum-Connes conjecture for groupoids*, Preprint, SFB 478 Geometrische Strukturen in der Mathematik, Mathematischen Instituts der Westfälischen Wilhelms-Universität, Münster, 1999.